

THE DARK SIDE OF GRAVITY

FREDERIC HENRY-COUANNIER

*CPPM, 163 Avenue De Luminy, Marseille 13009 France.
henry@cppm.in2p3.fr*

Newly added modifications in red in this version.

Adopting a non geometrical point of view for our space-time, we are led to an alternative theory of the order two and symmetric gravitational tensor field of GR. Considering our space-time to be flat and non dynamical, this field can no more be interpreted as its metric. The true metric is a globally Minkowskian background, a context which justifies a genuine rehabilitation of the global discrete space-time symmetries involved in the structure of the Lorentz group along with their 'problematic' representations: the negative energy and tachyonic ones. It turns out that about this background our gravitational field must appear in two discrete symmetry reversal conjugate forms, different but non independent symmetric order two tensors which must be treated on the same footing in our actions. The discrete symmetry reversal invariant actions, equations and their conjugate solutions are obtained. The conjugate form generates a Dark Side of Gravity. Indeed, matter living there can only interact antigravitationally with matter on our side. We show that stability is granted. In this Dark Gravity theory (DG), the new Schwarzschild solution in vacuum only starts to differ from that of General Relativity at the Post-Post-Newtonian order. DG does not violate WEP and though it violates SEP, the theory passes all present SEP tests except the Pioneer anomalous blue-shift, an SEP violating effect which is a natural outcome. No horizon (no Black Hole) arises in the Schwarzschild solution. A discontinuity of the gravitational field is propagating in the solar system. Depending on its present position, new gravitomagnetic effects well within the accuracy of the Gravity Probe B experiment might be observed in place of the GR frame dragging one. A spatially flat universe accelerated expansion phase is obtained without resorting to inflation nor a cosmological constant and the Big-Bang singularity is avoided. A helicity zero wave solution is obtained leading to the same binary pulsar decay rate as observed and predicted by GR. The context is also promising to help us elucidate several outstanding cosmological enigmas such as the flat galactic rotation curves.

Keywords: Negative energy; time reversal, tachyon.

1. Introduction

In special relativity, it is a well known and obvious result that the energy, being the time component of a four-vector must flip its sign under time reversal. It was also soon recognised however that negative energy objects lead to intolerable instabilities in our theories and should be discarded. In a quantum framework this is easily done thanks to the complex conjugation involved in the antiunitary time reversal operator of our Quantum Field Theories which avoids the generation of negative energies under time reversal. In Ref. 6 we nevertheless wanted to come back on the less conventional mathematical option e.g. that the time reversal operator be

unitary which not surprisingly reintroduces negative energy fields. We then noticed that, since there is no way to reverse a scalar field Hamiltonian sign under time reversal, it is impossible to reach a coherent description of unitary time reversal at least in a nongravitational framework.

We here propose a modified theory of gravity on flat space-time which extends, clarifies and explores in much more details some of our ideas already published in Ref. 56 to solve the issue at a classical level and show that it leads to a new understanding of time reversal and by the way 'the problematic' negative energy and tachyonic representations of the Lorentz group. The reader is referred to Ref. 6 for our detailed investigation of negative energies and time reversal restricted to non gravitational QFT. In the next section, we only remind the main conclusions of this study. An interesting analysis can also be found in Ref. 8.

2. Negative Energies and Motivations for an Alternative Theory of Gravity

Let us gather the main information we learned from our investigation in Ref. 6 of negative energies in Relativistic QFT indicating that the correct theoretical framework for handling them should be a modified GR.

- Theoretical Motivations

In second quantization, all relativistic field equations admit genuine negative energy field solutions creating and annihilating negative energy quanta. Unitary time reversal links these fields to the positive energy ones. With the unitary choice, usual for all other symmetries in physics, reversing time does not mean going backward in time anymore. Positive and negative energy fields vacuum divergences we encounter after second quantization are unsurprisingly found to be exactly opposite. The negative energy fields action must be maximised. However, there is no way to reach a coherent theory involving negative energies in a non-gravitational framework. Indeed, if positive and negative energy scalar fields are time reversal conjugate, also must be their Hamiltonian densities and actions. But for a scalar field, the Hamiltonian density is simply a sum of squared terms so there is obviously no way to make it flip its sign under time reversal, at least in a non gravitational context. However, when gravity comes into the game, $g_{\mu\nu}$ terms are expected in the Hamiltonian expression and their behaviour under time reversal might lead us to a solution to this issue. But this solution certainly cannot be found in GR because it is a geometric theory of space-time: global space-time symmetries and their associated Noether currents are lost and in particular, time reversal is no more the well defined symmetry it was in flat space-time. Therefore we must find a new theory of gravitation on a flat space-time background (we shall name it Dark Gravity or DG) in order to hopefully rehabilitate negative energy objects.

- Phenomenological Motivations

In a standalone mirror negative energy world which fields remain non coupled to our world positive energy fields, stability is insured and the behaviour of matter and radiation is as usual. Hence, it is just a matter of convention to define each one as a positive or negative energy world as was already noticed by Linde. Otherwise, if the two worlds are allowed to interact, both signs of the energy being involved in this interaction, a promising new phenomenology can be expected. Indeed, many outstanding enigmas seem to be suggesting that repelling gravity plays an important role in physics: flat galactic rotation curves, the flatness and acceleration (Ref. 41) of the universe, its voids...But negative energy particles never manifested themselves up to now in our detectors, also suggesting that a barrier is at work preventing the two worlds to interact except through gravity. The DG theory framework will involve by construction such a barrier.

The concordant cosmological SM is of course successful in its confrontation with as many precision observables as are the CMB, LSS, BAO and primordial elements abundances but this is at the price of simplicity since many added components and ideas of different kinds along with their free parameters enter into the game. These are dark energy, dark matter, inflation, some of them remaining badly understood and introducing new issues such as the fine tuning and coincidence problems. The impression that this construction is at least effective if not epicyclic has therefore recently motivated the research for alternative theories of gravity such as MOND (Ref. 35) and its relativistic extension TeVeS (Ref. 36). Part of our motivation for DG is similar.

- A Modified GR to Circumvent the Main Issues

A trivial cancellation between positive and negative vacuum divergences is not acceptable since the Casimir effect shows evidence for vacuum fluctuations. But in DG we will get cancellations of vacuum gravitational effects only.

Also, a generic catastrophic instability issue arises whenever positive and negative energy fields are allowed to interact. Restricting the stability issue to the modified gravity of DG, we shall show why this disastrous scenario is avoided.

Finally, even neglecting the former instability issue, allowing both positive and negative energy virtual bosons to propagate an interaction simply makes it vanish. Because electromagnetic, weak and strong interactions propagated by positive energy bosons will be isolated from the ones propagated by negative energy bosons, each being restricted to a different side of our gravitational barrier, the vanishing interaction issue remains a concern only for gravity in DG. However, the gravity interaction also will be treated very differently in DG than it is in GR so that this unpleasant feature, expected when gravitons of both energy signs can participate to the interaction, will also be avoided here.

3. Theories of Gravity on Flat space-time

Understanding negative energies as explained in the previous section is a strong motivation for looking for a flat spacetime theory of gravity but constructing a viable quantum theory of gravity is an even stronger one. Indeed, in the seventies, due to the increasing difficulties in trying to reach a coherent theory of quantum gravity, many theorists were led to the conviction that we might have to come back to a flat space-time theory of gravity and there were many attempts along this line. After all, because geometric considerations play no role in the derivation of GR which only rests on the general covariance requirement, it remains possible to adopt a non geometrical viewpoint in GR even if the gravitational field affects the space and time intervals we measure and even if it has the required mathematical properties to be considered as the genuine metric of space-time. Indeed, even in GR, we could still consider that the gravitational field has nothing to do with deformations of space-time which remains flat with true metric the globally Minkowskian $\eta_{\mu\nu}$. Just as in optical geometry, where we do not understand the curved trajectory of a light ray in a varying index medium as the effect of a deformed space-time but as the consequence of the interaction of light with the fields involved in this medium, just in the same way do we consider all the effects of gravity on our observables as due to mere interactions of our clocks and rods with the field $g_{\mu\nu}$, as any other field defined on a flat space time background described by $\eta_{\mu\nu}$.

However, as has been often pointed out, because $\eta_{\mu\nu}$ is absent from GR equations, there is no way to observationally distinguish the geometrical from the non geometrical point of view so that this issue remains one of metaphysics in the case of GR.

But, taking serious the flat space-time metric $\eta_{\mu\nu}$ makes it possible and even necessary to build different kinds of alternative theories of gravity. The so called multimetric theories exploit the fact that $g_{\mu\nu}$, if it can still be a metric topologically defined on our space-time is not considered as the metric of this space-time, the geometry defined by this metric having nothing to do with the geometrical properties of space-time. Then we are free to introduce as many fields of the $g_{\mu\nu}$ type as we want on a manifold without having to make appeal to extra-manifolds or extra-dimensions. Following this line of thinking, Petit (2 with references therein) has built a bimetric theory which associated phenomenology is very similar to the one we shall derive in DG. See also a review in Ref. 4 and references therein. In Rosen's (Ref. 30) approach, the true flat space-time metric $\eta_{\mu\nu}$ is explicitly introduced in the action. This theory is reviewed by C.M Will Ref. 21 and found to be ruled out by tests of the Strong Equivalence Principle (SEP) due to its background dependence. Because generically all background dependent theories are in trouble with the very constraining tests of SEP, the flat space-time (or background dependent) approach was progressively given up. DG is also a background dependent theory and it also violates SEP, though most such violations remain too small to be detectable thanks to fortunate compensations of the largest SEP violating terms in our field equations.

But LPI violation in space beyond the Post Newtonian order or LPI violation in time (Pioneer like effects) should eventually allow us to test DG against GR.

In the next section, we argue that not only the building of a theory involving both $g_{\mu\nu}$ and $\eta_{\mu\nu}$ is possible without conflicting with present observational Post-Newtonian constraints but moreover that our Dark Gravity theory is the minimal one that respects a new essential symmetry that went unnoticed in the 70's attempts.

4. Conjugate Worlds Gravitational Coupling

Now let us settle down the basic principles of DG. As usual for any order two tensor, symmetric and covariant gravitational field $g_{\mu\nu}$ we define its inverse, the contravariant $[g^{-1}]^{\mu\nu}$. But since $g_{\mu\nu}$ is not understood to be the metric of space-time, the latter is a contravariant tensor $\tilde{g}^{\mu\nu}$ but not the $g^{\mu\nu}$ tensor one would obtain by raising $g_{\mu\nu}$ indices using the true metric $\eta_{\mu\nu}$. Though the Ricci scalar one builds from $g_{\mu\nu}$ and $[g^{-1}]^{\mu\nu}$ is of course left unchanged by the pure renaming of $[g_{\mu\nu}]^{-1}$ into $\tilde{g}^{\mu\nu} = \eta^{\mu\rho}\eta^{\nu\sigma}\tilde{g}_{\rho\sigma}$, this rewriting reveals the existence of a second form of the field, still an order two symmetric covariant tensor $\tilde{g}_{\mu\nu}$ different but not independent from $g_{\mu\nu}$: this is $\tilde{g}_{\mu\nu} = \eta_{\mu\rho}\eta_{\nu\sigma}[g^{-1}]^{\rho\sigma} = [\eta^{\mu\rho}\eta^{\nu\sigma}g_{\rho\sigma}]^{-1}$. Thus the Janus gravitational field, like the Janus god, has two faces, $g_{\mu\nu}$ and $\tilde{g}_{\mu\nu}$ linked by the above manifestly covariant and background dependent relation. As the following simple picture makes it obvious, the two forms play perfectly equivalent roles and should be treated on the same footing in our actions if we don't want to artificially destroy the basic symmetry under their permutation (moving vertically inverses the tensors, diagonally raises or lowers their indices with $\eta_{\mu\nu}$).

$$[g^{-1}]^{\mu\nu} \quad [\tilde{g}^{-1}]^{\mu\nu}$$

$$g_{\mu\nu} \quad \tilde{g}_{\mu\nu}$$

Symmetrising the roles of $g_{\mu\nu}$ and $\tilde{g}_{\mu\nu}$ is performed by simply adding to the usual GR action, the similar action built from $\tilde{g}_{\mu\nu}$ and its inverse. Indeed, it is possible to have a connection compatible with $\tilde{g}_{\mu\nu}$ just in the same way as we had one compatible with $g_{\mu\nu}$ and exploit the mathematical apparatus of GR to constrain the general coordinate scalar built out of this form and its derivatives, another Ricci scalar. The theory that follows from symmetrising the roles of $g_{\mu\nu}$ and $\tilde{g}_{\mu\nu}$ is DG which turns out to be essentially the other option of a binary choice that must be done at the level of the conceptual foundations of a covariant theory of a symmetric order two tensor field: either the space-time is curved with metric $g_{\mu\nu}$ and we get GR, or it is flat with metric $\eta_{\mu\nu}$ and we get DG! GR actually just corresponds to the special case where g and η identify in which case, taking the inverse is equivalent to twice raising or lowering the indices with the space-time metric $g_{\mu\nu} = \eta_{\mu\nu}$. At last,

even if we did not assume from the beginning that $\eta_{\mu\nu}$ is the metric of space-time, in presence of the latter and a gravitational field $g_{\mu\nu}$, we automatically have the $g_{\mu\nu}$ and $\tilde{g}_{\mu\nu}$ forms with perfectly symmetric roles to play. Both being minimally coupled to matter and radiation fields, hence determining measured space and time intervals, could a priori be considered as the genuine metric of our space-time. However they are incompatible in general (they would describe a different geometry) so none of the two can be preferred so that it only remains $\eta_{\mu\nu}$ to play the role of the true space-time metric.

Since we are working on a globally flat space-time we will be free to use the flat space-time tools to build the Noether current associated with global symmetries and obtain a genuine energy momentum tensor which is not the GR energy-momentum pseudotensor. This was also Rosen's initial motivation for preferring flat space-time and many others followed him in this way (see Ref. 39–40 and references therein). We can also rehabilitate global discrete symmetries which are now well defined in a flat space-time framework along with their associated negative energy and tachyonic representations. We anticipate that we will have the good surprise to notice after solving our equations that the two faces of our Janus field solutions will be related by obvious discrete space-time symmetries. For instance, the $g_{\mu\nu}(t)$ and $\tilde{g}_{\mu\nu}(t)$ cosmological solutions will be found to be conjugate under time reversal followed by the more unusual space-time exchange symmetry which reverses the field signature (the most natural link between tachyons and bradyons): $g_{\mu\nu}(t) = -\tilde{g}_{\mu\nu}(-t)$ in the coordinate system where $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$.

Now let us try to deeper understand the meaning of $g_{\mu\nu}$ and $\tilde{g}_{\mu\nu}$. From Ref. 9 we learned that a discrete space-time symmetry, for instance time reversal, applying only to general coordinates, has no special role to play since it just belongs to the wide class of general coordinate transformations, and thus does not affect a scalar action. However, if the locally inertial coordinates $\xi^\alpha(x^\mu)$ can jump in a non-trivial way to another set of locally inertial coordinates $\tilde{\xi}^\alpha(x^\mu)$ at each space-time point:

$$\xi^\alpha(x^\mu) \rightarrow \tilde{\xi}^\alpha(x^\mu), \quad (1)$$

our action is not invariant anymore. Having two conjugate locally inertial coordinate systems at each space-time point, following the usual procedure, we of course have the associated two discrete symmetry reversal conjugate metric tensors:

$$g_{\mu\nu} = \eta_{\alpha\beta} \frac{\partial \xi^\alpha}{\partial x^\mu} \frac{\partial \xi^\beta}{\partial x^\nu}, \quad \tilde{g}_{\mu\nu} = \eta_{\alpha\beta} \frac{\partial \tilde{\xi}^\alpha}{\partial x^\mu} \frac{\partial \tilde{\xi}^\beta}{\partial x^\nu}. \quad (2)$$

(where the locally Minkowskian form of the fields, $\eta_{\alpha\beta} = \text{diag}(-1, +1, +1, +1)$ here has nothing to do with our global flat space-time metric of course) and the discrete space-time symmetry is only verified provided we add to the GR actions the ones built in the same way from $\tilde{g}_{\mu\nu}$. Then, we have to distinguish between those fields coupling to $\tilde{g}_{\mu\nu}$ and the others, coupling to $g_{\mu\nu}$. These matter and radiation fields of two mirror standard models respectively living in the two different forms of the gravity field never meet since they do not couple to each other. However, the relation between the conjugate forms, $\tilde{g}_{\mu\nu} = \eta_{\mu\rho} \eta_{\nu\sigma} [g^{-1}]^{\rho\sigma}$, allows these matter fields

to interact antigravitationally as we shall check later since a positive mass induced curvature of one form translates into a negative mass induced curvature of the conjugate form still over our flat space-time. But DG not only has two sectors, one of which being made of dark repelling matter from our side point of view. The introduction of an additional Einstein Hilbert action for the conjugate form also modifies the geometric side of the Einstein equation after exploiting $\tilde{g}_{\mu\nu} = \eta_{\mu\rho}\eta_{\nu\sigma}[g^{-1}]^{\rho\sigma}$ in order to eliminate $\tilde{g}_{\mu\nu}$. But this will have mainly PostPostNewtonian consequences, only non negligible in the cosmological and strong gravity domains.^a

In order to derive the fundamental equations of DG, solve them and check that we have a complete and acceptable theory (coherent and in good agreement with all observational data) we shall proceed in several steps: In step one, space-time permutation symmetries and discrete symmetries allow us to freeze various degrees of freedom and a priori identify only two allowed forms for the field in a privileged coordinate system where $\eta_{\mu\nu}$ reads $\text{diag}(-1, +1, +1, +1)$:

$$g_{\mu\nu} = \text{diag}(B, A, A, A) \implies \tilde{g}_{\mu\nu} = \text{diag}(B^{-1}, A^{-1}, A^{-1}, A^{-1}) \quad (3)$$

with either $B = -A$ or $B = -A^{-1}$. In step two we build the action sum of usual I_G and the conjugate one $\tilde{I}_G$ built in the same way from $\tilde{g}_{\mu\nu}$. The conjugate actions are separately general coordinate scalars and adding the two pieces is necessary to obtain a discrete symmetry reversal invariant total action. In step three, we vary together the unfrozen conjugate matrix elements, eliminate $\tilde{g}_{\mu\nu}$ thanks to the relation $\tilde{g}_{\mu\nu} = \eta_{\mu\rho}\eta_{\nu\sigma}[g^{-1}]^{\rho\sigma}$ and apply the extremum action principle to finally obtain our modified Einstein equation. In step four, we solve it and discuss our conjugate solutions in terms of discrete space-time symmetries. Exploring the phenomenological consequences of DG, we understand why those particles living in the conjugate world are seen from our world as negative mass particles. It is already clear why the gravitational barrier involved in DG explains the non detection of these particles and isolates the stability issue in the gravitational sector: these live on the other side

^a It was shown by Straumann in Ref. 26 that the building of a spin two $h_{\mu\nu}$ field theory over flat space-time automatically leads to identify a metric field and that its action turns out to be nothing but the Einstein-Hilbert action of GR. It follows that GR is the only theory of such spin two field perturbation over flat space-time, an argument which has been used to deny the validity of DG. The evidence that our action is eventually not the GR action after eliminating $\tilde{g}_{\mu\nu}$ simply tells us that DG is in no way the spin two theory over flat space-time Straumann started from. The deep reason why it is not is the gauge invariance requirement of the Straumann spin two theory, which is not meaningful in DG since it is not the general covariant theory of a single field, but of two separate fields $g_{\mu\nu}$ and $\eta_{\mu\nu}$. Thus DG is not invariant under an infinitesimal general coordinate transformation that only applies to $g_{\mu\nu}$ and not separately to $\eta_{\mu\nu}$, such general coordinate invariance condition being the one that usually translates into the gauge invariance requirement for $h_{\mu\nu}$ entering in $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ (the invariance of $\eta_{\mu\nu}$ under infinitesimal transformations was recognised as an important drawback of the perturbative treatment even in the context of GR: Ref. 66). By the way, having lost the usual gauge invariance of gravity in DG also puts under question a pillar of quantum field theory: gauge invariance as a building tool toward a theory of the electromagnetic or any other field. For completeness we mention that S. Hossenfelder (Ref. 44) has also recently initiated a study of anti-gravitation following probably the closest approach to the one followed here as far as we know.

of gravity preventing them from interacting weakly, strongly or electromagnetically with our side particles.

5. Space/Time Permutation and Discrete Symmetries

We carry out the program in four steps outlined at the end of the previous section, starting from step 1: in this section we will justify from global symmetry principles of flat space-time the only two allowed forms, $B = -A$ or $B = -A^{-1}$ in:

$$g_{\mu\nu} = \text{diag}(B, A, A, A) \implies \tilde{g}_{\mu\nu} = \text{diag}(B^{-1}, A^{-1}, A^{-1}, A^{-1}) \quad (4)$$

in a privileged coordinate system where $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$. These are not covariant but matrix relations a priori only valid in some particular frames. This is not a concern since we will keep working exclusively in such frames until our equations are solved for their tensor solutions which subsequently can easily be exported in whatever coordinate system we want. Our fundamental symmetry principle is that the form of $g_{\mu\nu}$ in these frames must first reflect the symmetrical roles of spatial coordinates. This requirement implies that $g_{\mu\nu}$ must be form invariant under matrix elements permutation of spatial indices which is enough to impose that for any spatial indices i, j : $g_{ti} = D$, $g_{ii} = A$ and $g_{i \neq j} = C$. But any rotation living invariant $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$ will destroy this form symmetry property unless $C=D=0$. Thus $g_{\mu\nu} = \text{diag}(B, A, A, A)$ in the privileged frames. Any Lorentz transformation (speed of light $c=1$) living invariant $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$ is also expected to destroy the $g_{\mu\nu} = \text{diag}(B, A, A, A)$ form. However, if we really take serious the flat space-time point of view, the observables should be defined with respect to an "exterior observer" able to measure with unrenormalised rods i.e absolute units in the sense that these are not affected by the gravitational field. But the speed of light as seen from such an "exterior observer" is no more the constant c . It is space-time (and side of the field) dependent through the effect of the gravitational field $g_{\mu\nu}$ and at each space-time point equals $c\sqrt{\frac{-B}{A}}$. It is clear that from such exterior observer point of view the true speed of light that enters the Lorentz transformations is the latter so that these Lorentz transformations apply differently at each space-time point, the appropriate point for studying special relativistic effects on a given object being at the object location. With this new understanding, Lorentz transformations, because they depend on the local value of the gravitational field, have nothing to do with the global symmetries of space-time. The latter should be restricted to homogeneity and isotropy i.e translation and rotation invariance. Anyway there is no conserved quantity associated with Lorentz boost in field theory so we loose nothing.

Of course, an actual observer is not exterior. Its rods are renormalised by the gravitational field so that he measures the constant speed of light c in any frame. Lorentz transformations applied by the actual observer keep invariant this c just as in General Relativity. But then, the fact that the two sides of the gravitational field take a diagonal form only in a particular Lorentz frame should not bother us: Lorentz

transformations don't have the same status as rotations or translations in DG. DG is obviously a theory with possible LLI violations and we will have to investigate them in details later. The important question is whether the LLI violating effects only arise in tests that were not already performed. We shall argue an affirmative answer in our section concerning DG gravitomagnetic predictions.

Let us stress the main differences between the procedures used in GR and DG to simplify the form of the gravitational field:

- In GR, isometries which can be expressed in the covariant language of Killing vectors allow to establish the existence of a particular coordinate system where a matrix relation is satisfied by the metric, for instance $g_{\mu\nu} = \text{diag}(B, A, A, A)$ in cartesian coordinates because it is believed that the metric must satisfy the isometries imposed by the symmetries of sources distribution. The procedure used in DG can in general be completely independent of these symmetries : a very strange feature which is only meaningful and tenable in the context of a theory having a background that can be exploited to define the preferred coordinate systems where the matrix relation is satisfied. As a consequence, even in a completely anisotropic source configuration and whatever their motions relative to each other and relative to our frame, our gravitational field is in the "isotropic form" in this privileged frame because this frame and nothing else imposes this form of the metric. Quite surprisingly, we will show that having enough privileged frames to avoid a conflict with constraints on the Lorentz violating parameters of the PN formalism, the predictions of DG can still agree with observations whatever the sources configuration.
- The two conjugate forms must have the same form (by the way, this implies also that they have the same isometries) and we can check that fortunately "the isotropic Cartesian form" $g_{\mu\nu} = \text{diag}(B, A, A, A)$ is also respected by the conjugate form in our privileged frame. This would not have been the case starting for instance from the Standard form ($g_{\mu\nu} = \text{diag}(B, A, r^2, r^2 \sin^2(\theta))$) usually used to derive the Schwarzschild solution in GR or even the isotropic form in polar coordinates ($g_{\mu\nu} = \text{diag}(B, A, Ar^2, Ar^2 \sin^2(\theta))$) since the inverse of a matrix element such as $g_{\phi\phi} = (r\sqrt{A(r)}\sin(\theta))^2$ leads to a strange $\tilde{g}_{\phi\phi} = \frac{1}{(\sqrt{A(r)}.r.\sin(\theta))^2}$ matrix element for $\tilde{g}_{\mu\nu}$ and it is easily checked, considering all its elements, that the latter field cannot also describe an isotropic solution.
- The procedure can efficiently reduce the number of degrees of freedom of $\tilde{g}_{\mu\nu}$ and $g_{\mu\nu}$ which is much a stronger constraint than in GR. Indeed, in some cases the off diagonal elements will not be varied at all (frozen to zero before varying the action) and the other elements will not be varied independently.^b

^bArguments against the freezing of fields elements before varying the actions have been given by

With the cartesian isotropic form condition for both forms

$$d\tau^2 = Bdt^2 + Ad\sigma^2, \quad d\tilde{\tau}^2 = \frac{1}{B}dt^2 + \frac{1}{A}d\sigma^2. \quad (5)$$

where $d\sigma^2 = dx^2 + dy^2 + dz^2$, we are left with the only two degrees of freedom A and B. We can now generalize to space-time the kind of arguments we have used to constrain the form of our field in the preferred frame by imposing a relation between the space-space A and the time-time matrix element B allowing to further reduce the number of independent elements to a single one. Then the equations of gravity might reduce to a single one with a single remaining degree of freedom to be varied. To do so we must identify a symmetry linking time and space coordinates. We know that the meaningful global symmetries on flat space-time should relate the fundamental representations of the Lorentz group i.e. tachyonic and bradyonic fields as well as positive and negative energy fields. The space/time symmetry reversing the signature provides a natural link between the tachyonic Lorentz group representation and the other ones according to the arguments given in Ref. 20. This leads us to suspect that even another conjugate form will be involved and should be introduced, the one with the flipped signature : $\hat{g}_{\mu\nu} = -g_{\mu\nu}$. Indeed, following the same method as before i.e. constraining the form of our field to be symmetric in the permutation of coordinates we would find the trivial B=A=C Euclidian form to be the only acceptable one because it manifestly involves time and space coordinates in a symmetrical way, whatever C(r,t) in $d\tau^2 = C(dt^2 + d\sigma^2)$. But if we also have the opposite form $\hat{g}_{\mu\nu} = -g_{\mu\nu}$ to exchange the role of A and B and restore in this way a kind of permutation symmetry between space and time we can certainly at least admit solutions with B=-A. Let us try to follow this method in an exhaustive way exploiting both conjugate forms $\hat{g}_{\mu\nu}$ and $\tilde{g}_{\mu\nu}$ to make our permutation symmetry work. **The most general expression is**

$$d\tau^2 = C \left(\frac{1}{D}dt^2 + Dd\sigma^2 \right), \quad d\tilde{\tau}^2 = \frac{1}{C} \left(Ddt^2 + \frac{1}{D}d\sigma^2 \right). \quad (6)$$

However D must be purely imaginary if we want a Minkowskian signature so we also need a purely imaginary C term to have a real field. Thus let us redefine D and C into iD and iC with the new D and C real. Introducing the opposite form

D.Deser and J.Franklin in Ref. 27 It is argued that in electromagnetism, isotropy a priori imposes that the magnetic field vanishes, however using this to freeze the magnetic field degrees of freedom before varying the action, one loses a fundamental equation which is the electromagnetism equivalent of the Birkhoff theorem: the vanishing of fields time derivatives in a spherically symmetric context. However this argument is only tenable provided there is no fundamental monopole in nature, since in presence of a monopole, B does not vanish and thus there is no reason to freeze it. But the inexistence of the monopole is certainly an a priori dangerous assumption so far. Moreover, our freezing of degrees of freedom does not follow from isometry requirements here as we have stressed. Ref. 27 also reminds us that freezing the off diagonal terms a priori in GR, one loses a fundamental equation from which the Birkhoff theorem is derived. But there is a priori no fundamental reason for the Birkhoff theorem to remain valid in DG (no need to protect this theory against radiation of scalar modes by an isotropically breathing source for instance).

$\hat{g}_{\mu\nu} = -g_{\mu\nu}$, we have eventually four conjugate forms $g_{\mu\nu}$, $\tilde{g}_{\mu\nu} = \eta_{\mu\rho}\eta_{\nu\sigma}(g^{-1})^{\rho\sigma}$, $\hat{g}_{\mu\nu} = -g_{\mu\nu}$, $\tilde{\hat{g}}_{\mu\nu} = \hat{g}_{\mu\nu}$ to hopefully describe conjugate positive/negative energy worlds as well as tachyonic/bradyonic worlds:

$$\begin{aligned} d\tau^2 &= C \left(\frac{1}{D} dt^2 - D d\sigma^2 \right), d\tilde{\tau}^2 = \frac{1}{C} \left(D dt^2 - \frac{1}{D} d\sigma^2 \right) \\ d\hat{\tau}^2 &= -C \left(\frac{1}{D} dt^2 - D d\sigma^2 \right), d\hat{\tilde{\tau}}^2 = -\frac{1}{C} \left(D dt^2 - \frac{1}{D} d\sigma^2 \right) \end{aligned}$$

only C and D both different than one breaks down the space / time permutation symmetry so that the symmetry is insured provided either C or D equal one (this is because in these particular cases the conjugate fields just exchange the roles of D and $-1/D$ or of C and $-C$) in which case the field must either be of the form $B = -A^{-1}$ or of the form $B = -A$. Thus, our new permutation symmetry principles allow us to accept only two kinds of real Minkowskian fields: the $B=-1/A$ and $B=-A$ forms. We can even understand the generation of such forms starting from a primordial trivial and most symmetric form of the kind $B=A=1$ which does not require us to introduce any other conjugate form to recover the space-time permutation symmetry. Then a continuous spontaneous, $\pi/2$ phase rotation of the form $B=1/A$, would transform this euclidian form into a couple of space/time conjugate imaginary opposite Minkowskian forms such that $B = 1/A = -A = \pm i$: Another spontaneous rotation of $B = -A$ would lead to the real η field. This one is striking as a common starting point for the subsequent dynamical (non spontaneous) evolution of the $B=-1/A$ and/or $B=-A$ fields.

6. The Conjugate Fields and their Variations

In the following sections we require that the field elements be real and a priori linked by $A = -B$ or $A = -B^{-1}$.

We postulate that $A = -B$ is such a strong symmetry that a single degree of freedom eventually remains. This means that A and B elements being related can no longer be varied independently. Therefore, given any tensor $R_{\mu\nu}$ and for real elements making use of $\frac{\delta g_{i'i'}}{\delta g_{ii}} = \frac{g_{i'i'}}{g_{ii}}$ and $\frac{\delta g_{tt}}{\delta g_{ii}} = \frac{g_{tt}}{g_{ii}}$ in case $B=-A$, a typical action variation will be proportional to a trace:

$$\delta(g^{-1})^{rr} R_{rr} + \delta(g^{-1})^{tt} R_{tt} + \delta(g^{-1})^{\theta\theta} R_{\theta\theta} + \delta(g^{-1})^{\phi\phi} R_{\phi\phi} = \frac{\delta(g^{-1})^{rr}}{(g^{-1})^{rr}} R$$

while in case $B=-1/A$, we will vary independently all field elements to get as many equations. We could speak of a weakly broken space-time global symmetry in this case, weakly because $B=-1/A$ does not eliminate degrees of freedom when we extremize the action but can be used to simplify the equations once we have them (just as we do in GR when isometries allow to work with a simplified form of the metric).

We also need the relation between the relative variations of the inverse conjugate forms needed to obtain the gravitational equation in term of the components of a single form of the field: $\frac{\delta \hat{g}_{xx}}{\hat{g}_{xx}} = -\frac{\delta g_{xx}}{g_{xx}}$. Recall that the opposite $\hat{g}_{\mu\nu}$ and $\tilde{\hat{g}}_{\mu\nu}$ and their

respective actions are also needed for $B=-A$. However, because the Einstein-Hilbert action of $-g$ is opposite to the Einstein-Hilbert Action of g , the $\hat{g}_{\mu\nu}$ and $\tilde{\hat{g}}_{\mu\nu}$ Einstein-Hilbert actions must come into the game with a negative sign i.e. with an opposite gravitational constant $-G$ as suggested in Ref. 97 to avoid a trivial cancellation of the total Einstein-Hilbert action and also to avoid stability issues. But then the new Einstein-Hilbert actions are identical to the ones we already have so we will assume these can be taken into account simply by a convenient redefinition of the factor G and it will not be necessary to involve explicitly these actions in the following.

It will also turn out that our solutions will always satisfy relations such as for example $\tilde{g}_{\mu\nu}(-t) = g_{\mu\nu}(t)$ in case of a background field: this will confirm that $\tilde{g}_{\mu\nu}$ is a time reversal conjugate of $g_{\mu\nu}$.

Eventually, for $B=-A$ we will have the single equation in vacuum :

$$\sqrt{g}R - \frac{1}{\sqrt{g}}R_{g \rightarrow 1/g} = 0$$

where R is the familiar Ricci scalar.

7. The DG Fundamental Equations in Vacuum

In this section we start from the rewritten proptime

$$d\tau^2 = -Bdt^2 + Ad\sigma^2$$

so that our previous section conditions: $B=-1/A$ and $B=-A$ translate into $B=1/A$, $B=A$. We can perform the computation in the more convenient and usual polar coordinates, since with the isotropic form the A and B entering in the field elements are the same in cartesian and polar ones. But we remind the reader that only B and A get reversed when passing from $g_{\mu\nu}$ to $\tilde{g}_{\mu\nu}$ of course. Moreover, this convenient polar coordinate system may be defined with respect to an arbitrary origin since our field does not necessarily privilege a particular center of isotropy.

We find for the Ricci tensor elements:

$$\begin{aligned} R_{rr} &= \frac{A''}{A} + \frac{B''}{2B} - \frac{B'^2}{4B^2} - \left(\frac{A'}{A}\right)^2 + \frac{A'}{Ar} - \frac{1}{4}\frac{A'}{A}\left(\frac{B'}{B}\right) + \left(-\frac{1}{2}\frac{\ddot{A}}{B} - \frac{1}{4}\frac{\dot{A}}{B}\left(\frac{\dot{A}}{A} - \frac{\dot{B}}{B}\right)\right) \\ R_{\theta\theta} &= \frac{3}{2}\frac{A'r}{A} + \frac{A''r^2}{2A} - \frac{A'^2r^2}{4A^2} + \frac{1}{2}\frac{B'}{B}\left(\frac{A'r^2}{2A} + r\right) + \left(-\frac{1}{2}\frac{\ddot{A}}{B} - \frac{1}{4}\frac{\dot{A}}{B}\left(\frac{\dot{A}}{A} - \frac{\dot{B}}{B}\right)\right)r^2 \\ R_{\phi\phi} &= \sin^2\theta R_{\theta\theta} \\ R_{tt} &= -\frac{1}{2}\frac{B''}{A} - \frac{1}{4}\frac{B'A'}{A^2} + \frac{1}{4}\frac{B'^2}{BA} - \frac{1}{A}\frac{B'}{r} + \frac{3}{2}\left(\frac{\ddot{A}}{A} - \frac{1}{2}\frac{\dot{A}}{A}\frac{\dot{A}}{A} - \frac{1}{2}\frac{\dot{B}}{B}\frac{\dot{A}}{A}\right) \end{aligned}$$

The sum of variations when $B=A$ involves the Ricci scalar:

$$R_{B=A} = \left(3\frac{A''}{A^2} - \frac{3}{2A}\left(\frac{A'}{A}\right)^2 + 6\frac{A'}{A^2r}\right) + \frac{3}{A}\left(-\frac{\ddot{A}}{A} + \frac{1}{2}\left(\frac{\dot{A}}{A}\right)^2\right)$$

The equation in vacuum when $B=A$ follows from

$$\sqrt{g}R - \frac{1}{\sqrt{g}}R_{g \rightarrow 1/g} = 0$$

giving:

$$3A \left(\frac{A''}{A} - \frac{1}{2} \left(\frac{A'}{A} \right)^2 + 2 \frac{A'}{Ar} - \frac{\ddot{A}}{A} + \frac{1}{2} \left(\frac{\dot{A}}{A} \right)^2 \right) - 3 \frac{1}{A} \left(-\frac{A''}{A} + \frac{3}{2} \left(\frac{A'}{A} \right)^2 - 2 \frac{A'}{Ar} + \frac{\ddot{A}}{A} - \frac{3}{2} \left(\frac{\dot{A}}{A} \right)^2 \right) = 0$$

For an homogeneous field $A(t)$ satisfies :

$$A \left(-\frac{\ddot{A}}{A} + \frac{1}{2} \frac{\dot{A}}{A} \left(\frac{\dot{A}}{A} \right) \right) - \frac{1}{A} \left(\frac{\ddot{A}}{A} - \frac{3}{2} \left(\frac{\dot{A}}{A} \right)^2 \right) = 0$$

admitting both time dependent as well as a stationary solutions.

For $B=1/A$, the time-time equation in vacuum follows from

$$\sqrt{g}G_{tt} - \frac{1}{\sqrt{g}}B^2G_{tt,g \rightarrow 1/g} = 0$$

giving:

$$\frac{1}{A} \left(2 \frac{A''}{A} - 2 \left(\frac{A'}{A} \right)^2 + 4 \frac{A'}{Ar} \right) + \frac{3}{4} \left(\frac{1}{A^2} - A^2 \right) \left(\frac{\dot{A}}{A} \right)^2 = 0$$

The space-time equation reads (B replaced by $1/A$)

$$A \left(\frac{\dot{A}'}{A} - \frac{1}{2} \frac{A'}{A} \frac{\dot{A}}{A} \right) + \frac{1}{A} \left(-\frac{\dot{A}'}{A} + \frac{3}{2} \frac{A'}{A} \frac{\dot{A}}{A} \right) = 0$$

and the space-space equations (B replaced by $1/A$):

$$A(A^2 \left(\frac{\ddot{A}}{A} + \frac{1}{4} \left(\frac{\dot{A}}{A} \right)^2 \right) - \frac{1}{A^2} \left(-\frac{\ddot{A}}{A} + \frac{9}{4} \left(\frac{\dot{A}}{A} \right)^2 \right)) = 0$$

Making the assumption that $\dot{A}$ does not vanish, we redefine $A = e^a$. Integrating as much as we can the three equations then read:

$$\dot{a}^2 \sinh(a) = \frac{4}{3} \Delta a$$

$$\dot{a}^2 \sinh(a) = f(t)$$

$$\dot{a}^2 \cosh^{5/4}(2a) = g(r)$$

but the last two for instance are not compatible unless $\dot{a} = 0$. Therefore the solution for the $B=1/A$ field is time independent and the only remaining equation reads $\Delta a = 0$

8. Summary

Now that we are in position to solve the DG fundamental equations, let us take sometime to first recapitulate the conceptual foundations from which these were derived and the main steps in this derivation to show that, though these look very unfamiliar for the GR expert, this derivation completely follows from basic principles which themselves are obvious in a flat space-time framework.

- Contemporary physics relies on symmetry principles. In a pure Special Relativity framework, time reversal symmetry is fundamental and generates negative energy objects solutions of all propagation equations. Quantum theory comes into the game with a binary mathematical choice for time reversal symmetry. Either it is Anti-Unitarily performed and may be we are allowed to neglect the negative energy fields solutions or it is Unitarily performed in which case the negative energy fields being regenerated by time reversal must be taken serious. Our starting point was the Unitary choice.
- The Hamiltonian of the most simple field, the scalar field, does not reverse under Unitary time reversal. This is a dead-end unless we extend our thought to gravity to investigate its time reversal behaviour.
- Time reversal is a global discrete symmetry which is only meaningful in a flat space-time and not in GR curved space-time. Thus we need an alternative theory of gravity involving at least the flat non dynamical background space-time $\eta_{\mu\nu}$ and the usual $g_{\mu\nu}$ of GR. For the sake of simplicity we first investigate a classical theory for these fields.
- But given $\eta_{\mu\nu}$ and $g_{\mu\nu}$, inevitably arises one (and only one) covariant symmetric order two tensor $\tilde{g}_{\mu\nu}$, depending on the latter and having exactly the same status as $g_{\mu\nu}$. So we must respect the symmetry under their permutation in the theory making them play equivalent roles in our equations.
- This is done by adding to GR action the same one built from $\tilde{g}_{\mu\nu}$. $\tilde{g}_{\mu\nu}$ is eliminated thanks to the relation that links it to $\eta_{\mu\nu}$ and $g_{\mu\nu}$ and the equations satisfied by $g_{\mu\nu}$ are derived from the least action principle in the coordinate system where $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$
- This frame also imposes global permutation symmetries to the fields which can only take the B=-A and B=-1/A forms that generate two cohabiting sectors of the theory.
- The DG equations are simplified by the compensating $g_{\mu\nu}$ and $\tilde{g}_{\mu\nu}$ terms as well as by the simple forms taken by the fields and a reduced number of degrees of freedom (one in case B=-A). Thus these are easily solved. Remains to be investigated the viability of our solutions, interpret them and understand the articulation between the B=-A and B=-1/A sectors before confronting the theory with various observables.

9. Russian Dolls Actions

The background metric provides a straightforward procedure to combine $[g^{B=-1/A}]_{\mu\nu}$ with $[g^{B=-A}]_{\mu\nu}$, once $[g^{B=-1/A}]_{\mu\nu}$ has been obtained from the resolution of the equation deriving from its standalone action, into a total single one $[g^{tot}]_{\mu\nu}$ the total field to which matter and radiation fields will actually couple in another action. $[g^{tot}]_{\mu\nu} = \frac{1}{2}([g^{B=-A}]_{\mu\rho}[g^{B=-1/A}]_{\nu\sigma} + [g^{B=-A}]_{\nu\rho}[g^{B=-1/A}]_{\mu\sigma})\eta^{\rho\sigma}$. The $[g^{B=-1/A}]_{\mu\nu}$ field involved in this combination will no more be dynamical in the new action since it has already played its dynamics in its own separate action. In that sense, we have what we might call Russian Dolls Actions. The procedure is also very powerful to combine various fields initially of the type $[g^{B=-1/A}]_{\mu\nu}$, may be one such field (and rest privileged frame) for each elementary source particle or small region of static fluid, the fields being combined after they have been exported to a common coordinate system of course. One can check that the procedure works well even in such non diagonal configuration of the combined fields. But we also want to keep as general as possible and keep open the possibility for extended sources and fewer privileged systems.

- The first actions we build are for the purely stationary $B(r)=-1/A(r)$ conjugate forms: each will describe the Schwarzschild gravity of DG for each perturbation in the cosmic mean density. Indeed we know that the Poisson equation we got can be solved to give the potential only for a field that vanishes sufficiently rapidly at infinity. The solution would be static in a homogeneous universe. Therefore this field only applies to perturbations. The action for all perturbations managed by a given preferred frame involves a non dynamical source action to be introduced in the following section. It is derived from the conserved energy momentum tensor we shall get from the second action. All $B=-1/A$ elementary fields, i.e one for each preferred frame are then Lorentz transformed to a common frame then combined in the total $B=-1/A$ field.
- The second action we build involves the $B=-A$ field for which we can solve for a background $B(t)=-A(t)$ solution and its $B(r,t)=-A(r,t)$ perturbations exchanging energy and momentum with the SM dynamical sources as usual but not the spectator here non dynamical background made of the total Schwarzschild type gravitational field.

A $B=-A$ wave perturbation can be generated about both conjugate sides of the background $B(t)=-A(t)$ field as well as about the background η in between them which is also a background possible solution. This will introduce an important extra factor three later when we come to compute the energy radiated by the decaying pulsar. Indeed, as we shall show, these solutions are waves different from GR ones but as successful as the latter to describe the quantitative decay of the binary pulsar.

10. B=-1/A, The New Schwarzschild Solution

- The non dynamical gravific action

The next step is to postulate our non dynamical source action. We know that for instance the total $T^{\mu\nu}$ energy momentum tensor has been obtained from the SM field action in the usual way and since the localized density perturbation we want to deal with here is a static perfect fluid, we have for the only non vanishing elements $T^{00} = -\rho(1+\Pi)/B_{tot} = -\rho(1+\Pi)/B_{ext}B$ and $T^{ij} = p\delta^{ij}/A_{tot} = p\delta^{ij}/A_{ext}A$ where B_{tot} and A_{tot} stand for the total $B_{tot} = -1/A_{tot}$ field the matter and radiation fields are propagating in and we have separated two contributions to the gravitational potential energy: an external one and the B=-1/A field we want to solve the equation for. It is usefull to define the scalars $\bar{p}$ and $\bar{\rho}$ which in the privileged coordinate system take the value

$$\bar{\rho} = \rho(1+\Pi)/B_{ext}; \bar{p} = p/A_{ext}$$

include internal as well as gravital potential energy in the external field but dont depend on A and B. This allows us to rewrite $T^{\mu\nu}$ in the usual way for a perfect fluid $T^{\mu\nu} = \bar{p}g^{-1\mu\nu} + (\bar{\rho} + \bar{p})U^\mu U^\nu$ with $U^\mu U^\nu g_{\mu\nu} = -1$. Again $g_{\mu\nu}$ here only involves the B=-1/A gravitational field we need to compute. The source action for our side fields then has to be

$$\int \sqrt{\eta} T^{\mu\nu} \eta_{\mu\nu} d^4x$$

for the integrand under a variation of $(g^{-1})^{\mu\nu}$ to become $\eta_{\nu\sigma} g_{\mu\rho} T^{\rho\sigma} \delta(g^{-1})^{\mu\nu}$.

To recover at least Newtonian gravity, solving $2\Delta a(r) = n\pi GM\delta(r)$ in case the source is a point mass M, must give $a(r)=-2U(r)$ where $U(r)$ is the adimentional Newtonian potential $U(r) = \frac{-MG}{r}$ solution of $\Delta U(r) = 4\pi GM\delta(r)$. Thus n equals -16 and our total Action S_{grav} for gravity and the gravific sources had to be

$$S_{grav} = \frac{1}{16\pi G} \int d^4x (\sqrt{g}R + \frac{1}{\sqrt{g}}R_{g \rightarrow 1/g}) + \int d^4x \sqrt{\eta} (T^{\mu\nu} + \tilde{T}^{\mu\nu}) \eta_{\mu\nu}$$

provided we have a perfect fluid at rest we can treat with a single privileged frame on both sides of the field.

The dynamics of our side SM fields does not take place in S_{grav} but in S_{SM} the familiar GR action describing how dynamical radiation and matter fields propagate in the total $g_{\mu\nu}$ field (exchanging between kinetic energy and gravitational potential energy), interact among each other but also exchange energy with the B=-A field. Similarly, the dynamics of the conjugate SM fields takes place in another $S_{\tilde{SM}}$ action involving $\tilde{g}_{\mu\nu}$ in place of $g_{\mu\nu}$. These $S_{\tilde{SM}}$ and S_{SM} are the second Russian Doll ones announced in the

previous section and they are obtained by the usual covariantisation procedure of the flat space time SM actions except that in these actions the Schwarzschild fields in turn are external and non dynamical having already played their dynamics in the first Russian Doll action S_{grav} . Therefore the bidirectional dialog between matter and $B=-1/A$ gravitation does not take place in a single total action as in GR and as will be the case for the $B=-A$ field but each direction is handled in an independent action for the $B=-1/A$ field. Let us write down the complete system of equations for this field. The time-time equation reads:

$$\left(2\frac{A''}{A} - 2\left(\frac{A'}{A}\right)^2 + 4\frac{A'}{Ar}\right) + \frac{3}{4}\left(\frac{1}{A^2} - A^2\right)\left(\frac{\dot{A}}{A}\right)^2 = -16\pi G(\bar{\rho}A - \frac{\bar{p}}{A})$$

The space-time equation reads:

$$A\left(\frac{\dot{A}'}{A} - \frac{1}{2}\frac{A'}{A}\frac{\dot{A}}{A}\right) - \frac{1}{A}\left(-\frac{\dot{A}'}{A} + \frac{3}{2}\frac{A'}{A}\frac{\dot{A}}{A}\right) = 0$$

and the space-space equations:

$$A^2\left(\frac{\ddot{A}}{A} + \frac{1}{4}\left(\frac{\dot{A}}{A}\right)^2\right) - \frac{1}{A^2}\left(-\frac{\ddot{A}}{A} + \frac{9}{4}\left(\frac{\dot{A}}{A}\right)^2\right) = -16\pi G\left(\frac{\bar{p}}{A} - \bar{p}A\right)$$

This system clearly shows that the gravific energy momentum tensor had to be defined as we did but with $\bar{p} = \bar{\bar{p}} = 0$ whatever p to impose that pressure should not source gravity either for an extended source such as a star or for the basic perfect fluid elementary source that we are considering as a good candidate model for the gravific elementary particle (this will be discussed below).

Then, as in vacuum, the space-time and space-space equations are trivially satisfied provided the field does not depend on time.

- The point mass M solution

Our static solution for a point mass source reads

$$A = e^{\frac{2MG}{r}} \approx 1 + 2\frac{MG}{r} + 2\frac{M^2G^2}{r^2} \quad (7)$$

$$B = -\frac{1}{A} = -e^{\frac{-2MG}{r}} \approx -1 + 2\frac{MG}{r} - 2\frac{M^2G^2}{r^2} + \frac{4}{3}\frac{M^3G^3}{r^3}. \quad (8)$$

different from the GR one though in good agreement up to Post-Newtonian order. No coordinate singularity arises in our frame and it is straightforward to check that this Schwarzschild solution involves no horizon. The conjugate forms can be transformed into one another through $r \rightarrow -r$ or $M \rightarrow -M$. Of course, this interpretation requires r to be defined as $r = \pm\sqrt{x^2 + y^2 + z^2}$ depending on the form we are living in. Thus, it is tempting to consider that the discrete symmetry reversal involved here reverses mass (gravific rest energy) and involves at least space reversal

$r \rightarrow -r$. Thus two conjugate fields (at r and $-r$) may appear antipodal with respect to $r=0$, the location of the point mass source.

Eventually it is important to notice that the poisson equation source term involves $\bar{\rho}A$ where A should affect importantly how the gravific mass varies inside matter but also the total integrated gravific Mass source as seen from outside matter because of the exponential behaviour in the strong field regime. In case of a very compact object this should contribute to generate a huge gravific mass very different from the gravific mass of the same quantity of matter spread over a larger volume. An interesting feature to understand the origin of the billion solar masses (this is a gravific mass) objects at the center of some galaxies.

- The PN solution

Adapting the PN formalism to DG, following the notation conventions of Ref. 21 we can systematically compare the obtained results with the corresponding ones in GR. The DG $B=-1/A$ field at PN order in the preferred frame reads:

$$\begin{aligned} g_{00} &= -1 + 2U - 2U^2 + 4\Phi_1 + 2\Phi_3 + 4\Phi_2 \\ g_{0j} &= 0 \\ g_{jk} &= (1 + 2U)\delta_{jk} \end{aligned}$$

In comparison, the GR PPN field at PN order reads:

$$\begin{aligned} g_{00} &= -1 + 2U - 2U^2 + 4\Phi_1 + 4\Phi_2 + 2\Phi_3 + 6\Phi_4 \\ g_{0j} &= -\frac{7}{2}V_j - \frac{1}{2}W_j \\ g_{jk} &= (1 + 2U)\delta_{jk} \end{aligned}$$

where

$$U = (x, t) = \int \frac{\rho(x', t)}{|x - x'|} d^3x'$$

and Φ_4 , V_j , W_j terms are respectively pressure and momentum source terms that appear in GR but not in DG in the preferred frame while Φ_1 , Φ_2 , Φ_3 are kinetic energy, gravitational potential energy and internal (non gravitational potentials) energy terms. The explicit expressions for these terms can be found in Ref. 21 p95. The pressure contribution Φ_4 can be removed in DG in the preferred frame simply by postulating that the microscopic motions of the fluid particles do not contribute to gravity through pressure i.e by defining $\bar{p} = 0$ and $\tilde{p} = 0$ in place of the previous section definitions. Then we still have the important result we already had in vacuum: the $B=-1/A$ field is stationary. Φ_4 could not be isolately measured up to now which according to Ref. 28, is due to an exact compensation between pressure and gravitational potential energy of a star in a static configuration. Therefore, DG is not yet constrained in this sector. We see that in our framework, the momentum and pressure source terms can be avoided for an elementary field i.e. the field which contributions to the source are managed within a

single privileged frame. This is a trivial result in case the elementary field is the field of a single mass source at rest in the privileged frame it defines: its rest frame. In this case the static picture becomes the most natural one so the reader might wonder why we do not specialize from the beginning to this most natural and trivial scenario. The answer is that we want to keep open the possibility of making frame dragging predictions different from GR ones in some cases (in a forthcoming section).

Since the elementary field, before it is combined with others, is always independent of its sources momentum and pressure, in the privileged frame only the total energy (including all potential energies) can source our $B = -1/A$ gravity.

In case there is a preferred frame associated with each particle, pressure terms should eventually arise as the result of combining all the elementary fields produced by all elementary sources and the theory will eventually involve pressure gravific effects in the total gravitational field. However if an extended fluid gravific effect were to depend on a single privileged frame we would expect no pressure contribution. In this case, a possible way to test DG against GR would be an evidence for the active gravific pressure in a highly non equilibrium situation such as that of a Supernova.

Because the Φ_3 and Φ_2 potential energy terms are the same at PN order in DG and GR, no LLI nor LPI violation effects associated with these terms are expected in DG at PN order.

In a previous version of this article the kinetic Φ_1 was absent so all kinetic terms were absent in the DG solution which meant another departure from GR predictions. However we have just learned that the mass of the proton was recently successfully (with good precision) computed and is the total rest energy mostly determined by internal energies of the partons (QCD terms, and kinetic terms), the quark masses contribution remaining very small. Since the equality of gravific and inertial mass is well tested for baryonic matter, it would be dangerous to forget the Φ_1 contribution to the gravific mass in DG. However since we were not informed about the exact actual contribution of kinetic terms to the proton mass, we cannot exclude that Φ_1 be absent in DG.

Remains the absence of the momentum source terms which possible consequences will be explored in the next section. The Strong Equivalence Principle, as in any background dependent theory is menaced (see in Ref. 21 the discussion of Rosen's Ref. 30 bimetric theory for instance). However it seems that our framework successfully passes all present PN tests so far.

- The exact solution

In our privileged coordinate system, the elements read $-1/A = B = e^{2U(r)}$ (notice by the way that $-1/B = A$ is only satisfied in the standard coordinate system in GR, not in the isotropic one), where $U(r)$ is simply the total additive Newtonian gravitational potential generated by the mass distribution.

For a set of masses M_i , $U = -\sum_i \frac{2M_i G}{r_i}$. This is our "elementary" solution and we explained earlier how the total $B=-1/A$ field might be obtained as a superposition of such elementary Russian Doll actions $B=-1/A$ solutions, each corresponding to a privileged frame. Of course the combination must be performed after Lorentz transforming the elementary fields in a common coordinate system thus will include the corresponding kinetics effects of the sources. The theory is of course relativistic: in any other frame both the gravitational field and the source tensor will in general loose these simplest forms. For instance the gravitational field will loose its diagonal form and will have momentum and kinetic energy components.

- The gravific source Action

The physical mechanism behind the generation of our gravific action would be the following: we postulate the existence of a network of huge masses with alternate positive and negative values in vacuum at the microscopic scale. The strong anti-gravitational interaction between each mass and its neighbours stabilizes the network. At the locations of these mass does the creation and annihilation of spherical waves of QFT take place. In particular, the network is responsible for the QM wave collapse. Only at the location of these masses is the gravitational field generated when various waves are created and annihilated at these points. The huge masses also generate their own gravity but their compensating fields result in undetectable gravitational effects at the macroscopic scale.

It is understood that any incident wave packet can be trapped in the deep well of one of the network masses where it momentarily has the rest energy M (standing wave) before being reemitted after some delay with the same kinetic energy as initially. In between points of the network the fields propagate but never generate any gravitational field. The hope is to eventually provide a mechanism for the generation of a gravific mass which would simply be the total energy momentarily trapped in the well, even for a massless wave propagating at the speed of light in between the nodes of the network. Another interesting possibility to be explored is the following: Even a massless field far away from any discontinuity might acquire a mass (rest energy) in the vicinity of a discontinuity of the gravitationnal field (we shall encounter these discontinuities later when we study the $B=-A$ field). In principle the time coordinate can rotate in the complex plane at a discontinuity and we saw that the conjugate forms of the field can be transformed into each other by such rotation. But if such rotation can start in the vicinity of the discontinuity a well known extra dimensionnal effect could take place there: the momentum lost in the generated extra dimension would be seen as a mass from our four dimensionnal point of view. In this way the massless field may acquire a mass in the vicinity of the discontinuity and then be trapped in the volume enclosed by such discontinuity (while field remaining

massless would not be sensitive to $B=-A$ field discontinuities and could not be trapped). It would be trapped in the form of a system of standing waves by the discontinuity surrounding the deep potential well generated by any mass in the vacuum network. This is a possible way to generate gravific rest energy (mass) but only at the network Masses.

By the way this mechanism could also generate a tiny photon mass which would even allow the radiation pressure to contribute to the gravitational field after Lorentz exporting all elementary fields generated in each photon restframe in a common frame. This is an interesting possibility though not the DG most likely scenario we described earlier: no radiation pressure contribution to the gravitational field.

The last section will be devoted to deeper speculations following this line of thinking.

- Stability

The equation does not admit any wave solution. So the solution is unpropagated and instantaneous in our frame and there is no force carrier. Usually stability is menaced in a theory when there are interacting degrees of freedom with both positive and negative energy states opening an infinite phase-space for decay. Once the DG equations are obtained, the very brutal elimination of all but one degrees of freedom, $B=-1/A$, with direct consequence from our equations that it cannot depend on time in vacuum trivially insures stability of this sector of the theory.

One may wonder how the solution could remain static in case the sources were not static. First the issue seems real only when we want to describe extended source distributions for an elementary source is always static in its restframe. But even for the extended case our solution is only static in the sense that it has no explicit time dependency. Of course the set of variables r_i entering in our solution can change in time when the sources labelled by i are in motion.

The phenomenology is simple: masses living in the same form attract each other. Masses living in different forms repel each other, as if the mass living in the conjugate form contributes as a negative mass source from our side point of view. This is the same phenomenological stability as in the bimetric theory of JP Petit (Ref. 1 and 2 with references therein). This phenomenological stability is interesting to notice. Indeed, in GR we would naively expect a negative energy object to be attracted by a positive energy object, the latter being repelled by the former. They then would accelerate together for ever. Here such kind of phenomenological instability is also avoided since masses living in different forms just repel each other. Yet, from the point of view of each form, this is really the interaction between a positive mass and a negative mass.

- Causality

The proper time defined in association with the $B=-1/A$ field can be fac-

torized:

$$d\tau^2 = A(-dt^2 + \frac{1}{A^2}d\sigma^2)$$

Let us follow the point of view of the background spacetime metrics which unrenormalised rods would allow us to measure a light speed different from c . The A element affects the local speed of light seen by such fictitious observer $c_{local} = ca$ where c stands for the fundamental constant appearing everywhere else in our fundamental equations and $a = \sqrt{A}$ while in the conjugate form, the local speed of light the same observer would measure is $\tilde{c}_{local} = c/a$ greater than $c_{local} = ca$ for instance in the vicinity of a Mass source at rest on our side. In this case the conjugate light-cone is wider than our side lightcone. This remains true from the genuine observer point of view on our side and means that the speed of light on the conjugate side also appears to him greater than c he measures on his side. This possibility of faster than c information transfer in a special relativistic framework is well known to generate causality issues: some observers might see these informations propagating backward in time.

Having an instantaneous interaction in a theory raises an even more serious causality issue. Instantaneity can only be valid in one Lorentz frame. If the instantaneity frame is the restframe of the emitter, many emitters in motion relative to each other define the same number of different instantaneity frames (this is the case when we have many privileged frames, one per individual source) which makes it possible for a signal exchanged from A to B in relative motion, then back from B to A , to arrive in the past of A original emission.

Of course, the issue would disappear in case of a unic frame of instantaneity whatever the emitters motion relative to this frame. Indeed A could at most instantaneously send a signal to himself via B , never in its proper past, this remaining valid in any other frame: $\Delta t_{AB} + \Delta t_{BA} \geq 0$. But a single cosmic preferred frame would have produced large and easily detectable LLI violating effects in many solar system tests (see our next section devoted to gravitomagnetism).

Backward in time propagation of information in DG thus appears a real possibility and we shall take it serious. First, after all, we know from various tests of quantum non locality that there actually exists influences able to propagate faster than light (even instantaneously) even if as is usually believed, these cannot be used to transfer information. Secondly, propagation of informations from the future to the past is not a concern provided the whole story is coherent i.e. the events induced by such transfer of information will not affect the future events differently than what is already predetermined : indeed the information received from the future means that this future is already written and cannot be modified. Actually, it is only

when consciousness and in particular free will come into the game that the causality issue arises since it's difficult to see what would forbid this free will to act differently than what is already known from the written future. So this kind of issue is most probably mainly related to our fundamental ignorance of the deep nature of consciousness and may be also of what decides the wave function collapse in QM. For instance the collapse may not be complete in some cases allowing the separation of two parallel worlds if a free will decided to modify the course of the events: instead of such modification a distinct parallel future would be generated.

- WEPR

The Weak Equivalence Principle is obviously not menaced if once the field solution is established, matter and radiation have to follow its "geodesics" as in GR since the actions S_{SM} and $S_{S\bar{M}}$ are the same as in GR except for the presence of the B=-A field we shall study later on.

- Energy of the Gravitational Field

Most components of the energy momentum tensor $t^{\mu\nu}$ of the B=-1/A gravitational field, a Noether current computed thanks to the global translational invariance of the action, trivially vanish for the B=-1/A solution since this field is time independent. The Lagrangian

$$L = \frac{1}{16\pi G} \left(-2 \left(\frac{A'}{A} \right)^2 + 3A^2 \left(-\frac{\ddot{A}}{A} - \frac{1}{2} \left(\frac{\dot{A}}{A} \right)^2 \right) + 3 \frac{1}{A^2} \left(\frac{\ddot{A}}{A} - \frac{3}{2} \left(\frac{\dot{A}}{A} \right)^2 \right) \right)$$

has second derivatives so the expression of the conserved Noether current is a little bit less trivial than the usual one:

$$t_{\mu\nu} = \frac{\partial L}{\partial(\partial^\mu A)} \partial_\nu A - \eta_{\mu\nu} L - \partial^\rho \left(\frac{\partial L}{\partial(\partial^\mu \partial^\rho A)} \right) \partial_\nu A + \frac{\partial L}{\partial(\partial^\mu \partial^\rho A)} \partial_\nu \partial^\rho A$$

We could obtain this simple expression exploiting the fact that our Lagrangian only depends on second derivatives $\partial^\mu \partial^\rho A$ of a single coordinate ($\mu = \rho$). We find

$$t_{00} = -\frac{\partial L}{\partial \dot{A}} \dot{A} + L + \frac{\partial}{\partial t} \left(\frac{\partial L}{\partial \ddot{A}} \right) \dot{A} - \frac{\partial L}{\partial \ddot{A}} \ddot{A}$$

but all time derivatives vanish so $t_{00} = L = \frac{1}{-16\pi G} (2a'^2)$.

- Rotating Frames

An elementary $g_{\mu\nu}$ (B=-1/A) transforms as usual under a transformation from the privileged frame to any rotating frame. In the former, we already interpreted the expression of the proper time:

$$d\tau^2 = A(-dt^2 + \frac{1}{A^2} d\sigma^2)$$

On a disk uniformly rotating at $z=0$ about any z -axis with angular velocity ω the motion of a point P at distance ρ from the axis, allows to define a

rotating polar frame where the P polar coordinates $\rho, \theta, 0, t$ can be related to x, y, z, t through

$$x = \rho \cos(\omega t + \theta), y = \rho \sin(\omega t + \theta), z = 0$$

In the rotating frame, the proper time now reads:

$$d\tau^2 = -\left(A - \frac{\omega^2 \rho^2}{A}\right) dt^2 + \frac{\rho^2 d\theta^2}{A} + 2 \frac{\omega \rho^2}{A} dt d\theta$$

All special relativity effects are of course the same in GR and DG but the interpretation of these effects from the outside observer (with unrenormalized rods) point of view is worth describing in DG: For a clock at rest ($d\theta = 0$) in this frame, $d\tau^2 = -A\left(1 - \frac{\omega^2 \rho^2}{A^2}\right) dt^2$, and we see that the "speed of time" is modulated in a rotating frame by an extra factor $\left(1 - \frac{\omega^2 \rho^2}{A^2}\right)^{-1/2}$ relative to its $A^{-1/2}$ behaviour in the frame we started from. For a light signal, $d\tau = 0$ and the speed of light is given by solving the previous equation for dt to get $\frac{\rho d\theta}{dt} = A \frac{1 - \frac{\omega^2 \rho^2}{A^2}}{1 \pm \frac{\omega \rho}{A}}$ with an extra factor $\frac{1 - \frac{\omega^2 \rho^2}{A^2}}{1 \pm \frac{\omega \rho}{A}}$ relative to its A behaviour in the frame we started from. The speed of light thus appears anisotropic depending on light running clockwise or counterclockwise in the rotating frame. In GR, there is no such exterior background point of view and the anisotropic speed of light interpretation is not meaningful: the Special Relativity Sagnac effect is obtained because space and time intervals are deformed in the rotating frame.

- **Speed of Gravity**

Recently, Kopeikin suggested in Ref. 29 that the time delay of light from a quasar as the light passed by the Jupiter planet could be used to measure the finite speed of gravity. However, the analysis of the light propagation in the Jupiter's rest frame makes it obvious that the speed of gravity is irrelevant in this frame (Ref. 34) so that such measurement represents no more than a test of LLI. Up to now there thus exists no evidence for the finite speed of the gravitational interaction.

11. The total $B=-1/A$ field and cosmology

The total $B=-1/A$ field obtained by combining all elementary such fields cannot drive a global evolution because there is no solution of the poisson equation for an homogeneous source distribution. Thus our universe would be completely static without the contribution of the $B(t)=-A(t)$ cosmological background.

A void on our side or a concentration of matter on the conjugate side will both have a repelling gravitationnal effect on our side and attractive on the conjugate side so that such voids will appear to expand relative to their environment and reach a more perfect vacuum than in GR thanks to the conjugate structure that soon nonlinearly grows at the same time from the initial perturbations.

When this globally static picture is complemented by the effect of the field $B(t)=-A(t)$ all perturbation scales will be submitted to cosmological effects in exactly the same way (very different from what happens in GR).

12. $B=-1/A$, Gravitomagnetism

We now explore the gravitomagnetic sector of the theory. The main new DG prediction of this section will be the absence of frame-dragging effects below the scale encompassing all sources entering in the same "elementary $B=-1/A$ " field. If this scale just encompasses our earth, its rotation will have no effect on GP-B gyroscope axis. We follow the PPN formalism method as much as possible.

In a chosen PPN coordinate system moving at velocity $\vec{w}$ relative to our privileged frame, we can get the Lorentz transformed g_{0i} element to Post-Newtonian order for a point mass source m . We get

$$g_{0i} = -4w_i \frac{m}{r}$$

at the Post Newtonian order. Considering a set of such sources will result in many such w dependent terms in the field elements while the expression in GR of the same elements would also involve speeds of the sources relative to each other leading to sources angular momenta contribution in the PPN system (Ref. 21, p104).

Since $\vec{w}$ may be written as $-\vec{w} = \vec{w}_{PF} = \vec{w}_{Source} + \vec{w}_{PF/Source}$, g_{0i} involve a preferred frame term depending on the source velocity relative to the preferred frame where the field was diagonal and satisfied $B=-1/A$.

As a result, the precession of a gyroscope's spin axis $\vec{S}$ relative to distant stars as the gyroscope orbits the earth is given by (Ref. 21 , p208):

$$\frac{d\vec{S}}{d\tau} = \vec{\Omega} \times \vec{S}, \vec{\Omega} = -\frac{1}{2}\vec{\nabla} \times \vec{g} + \frac{3}{2}\vec{v}_{GPB} \times \vec{\nabla} \left(\frac{m}{r} \right)$$

the components of the 3-vector $\vec{g}$ being the g_{0i} . Explicitely,

$$\vec{\Omega} = -2\vec{w} \times \vec{\nabla} \left(\frac{m}{r} \right) + \frac{3}{2}\vec{v}_{GPB} \times \vec{\nabla} \left(\frac{m}{r} \right)$$

where $-\vec{w} = \vec{w}_{Source} + \vec{w}_{PF/Source}$. This yields

$$\vec{\Omega} = \vec{\Omega}_{\text{geodetic}} + \vec{\Omega}_{\text{PF}}$$

where in addition to the geodetic precession also expected in GR,

$$\begin{aligned} \vec{\Omega}_{\text{geodetic}} = & +2\vec{w}_{Source} \times \vec{\nabla} \left(\frac{m}{r} \right) + \frac{3}{2}\vec{v}_{GPB} \times \vec{\nabla} \left(\frac{m}{r} \right) = \\ & \frac{3}{2}\vec{v}_{GPB/source} \times \vec{\nabla} \left(\frac{m}{r} \right) + \frac{7}{2}\vec{w}_{Source} \times \vec{\nabla} \left(\frac{m}{r} \right) \end{aligned}$$

which last anomalous term (depending on the arbitrary PPN system) may be dropped since it does not appear in the truly measurable quantity (Ref. 21 , p211), a preferred frame like effect arises in $\vec{\Omega}_{\text{PF}} = -2\vec{w}_{Source/PF} \times \vec{\nabla} \left(\frac{m}{r} \right)$. Hence, we

have a Preferred frame Post Newtonian effect that comes in place of the Lense-Thirring precession or “the dragging of inertial frames” usually interpreted as a genuine coupling in GR between the spins of the earth and gyroscope. Indeed, we are free to choose the same w_{Source} , for instance here the extended body center of mass speed, for any elementary point mass m entering in the composition of this body, then factorize this speed when summing over all point masses to get the GR geodetic terms plus another term independent of the body angular momentum. By the way, the highly non trivial issue of deviation from geodesics for a spinning body in general relativity is also avoided if the body angular momentum does not source gravity.

It is instructive to compare this preferred frame effect to the preferred frame effect that arises in the Parametrized Post Newtonian formalism:

$$\vec{\Omega}_{PF} = \frac{1}{4}\alpha_1 \vec{w}_{PF} \times \vec{\nabla} \left(\frac{m}{r} \right)$$

Following Ref. 21, p209, for an earth orbiting satellite, the dominant effect comes from the solar term leading to a periodic angular precession with a one year period, with amplitude:

$$\delta\theta_{PF} \leq 5.10^{-3''}\alpha_1$$

thus completely negligible according to the PPN formalism given the experimental limit $\alpha_1 < 4.10^{-4}$.

Because in this general perspective DG just appears as an $\alpha_1 = -8$ (it is also straightforward to show that $\alpha_3 = -4$) theory one should wonder why we still take serious the possibility of detecting these preferred frame effects with GP-B and also in other classical tests of GR which up to now were able to constrain very strongly the α_i parameters. The answer is simply that we expect the preferred frames to depend on local changing conditions such as the position of various propagating discontinuities delimitating several domains with different $B(t)=-A(t)$ evolutions. Since we are able to explain the Pioneer anomaly (more details in the next section) relying on such discontinuity in the solar system, we suspect that various preferred frame effects may appear even in famous classical tests such as the secular change in perihelion position and geophysical tests where they did not show up before. Remember that the extreme case in DG in which we would have each source mass defining its own preferred frame would cancel all such preferred frame effects. Another possibility is that some important discoveries have been falsified or hidden by scientists working for military black programs.

Our effect is mainly determined by the solar term (in which the source is the sun) and has an amplitude $\delta\theta_{PF}$ approaching $0.02''$ for a preferred frame attached to the center of our galaxy and even greater for a preferred frame attached to the CMB, if the CMB actually follows the global motion of our present universe clusters (which might not be the case according to recent studies Ref. 93). In case the preferred frame is attached to the sun, the dominant term obviously vanishes while it yields a small effect comparable to the sun geodetic effect in case the preferred frame is

linked to the earth. The subdominant term is the earth (its source is the earth) one which vanishes in case the earth defines the preferred frame. If the sun defines the preferred frame, it produces a small periodic drift of the gyros with a GPB orbit period modulated by a term depending on the velocity of earth relative to the sun. If the center of our galaxy defines the preferred frame or in case of any more global frame an extra periodic term (with still the GPB orbital period) adds up depending on the velocity of the sun relative to this frame. A GPB gyro West Est (counted positive from West to Est) drift is illustrated over one year for the case of a preferred frame attached to the center of our galaxy in Figure 3, showing the dominant one year periodic drift and the two subdominant 1.6 hour periodic ones. $t=0$ is the spring equinox time. The global quasi linear behaviour on a limited period of up to six months around $t=0$ might well mimic the GR frame-dragging but the ambiguity must disappear once the full year data are considered. The mathematica plot does not allow to resolve the rapid 1.6 hour periodic oscillations and one should not pay attention to the seemingly chaotic behaviour of the curve but only to the smoothly varying upper and lower limits.

The Preferred frame term does not depend on the particular chosen working PN coordinate system but on the speed of our source relative to a privileged frame $\vec{w}_{Source/PF}$ in DG! Had the chosen PN coordinate system been the preferred one itself, the effect would have been there and could have been interpreted as a pure geodetic effect component depending on the speed of the source entering in a total geodetic effect depending on the speed of the test mass in this frame rather than the more usual speed of the test mass relative to the source. Let us notice that such effect is well reachable given the experimental accuracy ($5.10^{-4''}/year$) of the Gravity Probe B experiment designed to measure for the first time gravitomagnetism isolated from other Post-Newtonian effects. On a short period, it would even give a contribution on both North-South and Est-West directions slightly but significantly shifting the expected geodetic effect, and mimicking a frame-dragging like effect but would disappear integrated on a complete year.

Our conclusion is that if the preferred frame is a local one attached to the earth, the new DG prediction is the absence of frame-dragging, the rotating earth has no effect on a GP-B gyroscope axis. Moreover we predict the same geodetic effect as in GR. If the preferred frame is a global one (CMB frame) then we have the additional effect described above. But since this is a one year periodic oscillation, it does not contribute to the integrated effect. So the integrated prediction is the same but the drift is not a steady one which may provide an additional signature.

Let us repeat that in the very extreme and improbable limit where we would have as many preferred frames as elementary particles, our predictions would tend to the GR one: the same frame dragging and geodetic effects.

For the time being, Lunar Laser Ranging did not measure frame-dragging effects although there were recent claims by Ciufolini that frame-dragging effects due to the rotating earth have been already evidenced at the 10 percent level of accuracy

in an analysis exploiting the LAGEOS satellites data and indirectly in the binary pulsar PSR 1913+16, these are complicated analysis with undedicated data which sources of systematical effect may not be under total control so that it is more commonly believed that only the dedicated experiment GP-B measurement will hopefully soon give us the final answer. Moreover, the kind of preferred frame effect described above might also mimick a frame-dragging.

Another interesting possibility is the following: We shall show later that DG naturally predicts the occurrence of discontinuities of the background B=-A gravitational field allowing to quantitatively explain the Pioneer anomaly. Such discontinuities are propagating in the solar system. Therefore it is natural to suspect that these discontinuities of the gravitational field are delimiting spatial volumes and that each such volume is the domain of validity of a preferred frame. For instance, if at a given time a discontinuity was encompassing the earth and GP-B orbit, the local earth preferred frame did not allow any significant preferred frame effect to be detected. If more recently, the discontinuity has drifted and now occupies an equipotential surrounding the earth at an altitude lower than the GP-B orbit then GP-B is no more in the domain of validity of the earth preferred frame but rather of a sun or more global preferred frame in which case we might see the associated preferred frame effects described above. If the crossing of the discontinuity took place during the GP-B data taking then GP-B gyros would may be manifest the succession of the two regimes.

13. B=-A, the Cosmological Field

Our cosmological gravitational field only depends on time. By the way, even if we did not impose the B=-A condition, the only possible coordinate system with both forms spatially homogeneous and isotropic is a spatially flat one. In the global preferred coordinate system where $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$, a couple of time reversal conjugate purely time dependent solutions can be derived from the couple of conjugate actions. The existence of a time reversal conjugate universe was also suggested a long time ago in Ref. 15. For the single degree of freedom A(t) we were led to the equation :

$$3A \left(-\frac{\ddot{A}}{A} + \frac{1}{2} \left(\frac{\dot{A}}{A} \right)^2 \right) - \frac{3}{A} \left(\frac{\ddot{A}}{A} - \frac{3}{2} \left(\frac{\dot{A}}{A} \right)^2 \right) = n\pi G(A^2(\rho - 3p) - \frac{1}{A^2}(\tilde{\rho} - 3\tilde{p}))$$

The right hand side perfect fluid source term is just $\sqrt{g}T - \frac{1}{\sqrt{g}}T_{g \Rightarrow 1/g}$ obtained by the extremization of the usual SM and $\tilde{S}\tilde{M}$ actions where the fields are dynamical except the B=-1/A fields obtained in a previous section.

The covariant energy momentum conservation equations deriving from the SM actions can be written on both side: $T_{;\nu}^{\mu\nu} = 0$ and $\tilde{T}_{;\nu}^{\mu\nu} = 0$. They result in the evolution laws $\tilde{\rho} = 3\tilde{p} \propto 1/\tilde{A}^2$ and $\rho = 3p \propto 1/A^2$ in the radiative eras while $\tilde{\rho}$ and $\tilde{p}$ vary as $\propto 1/\tilde{A}^{3/2}$ and ρ and p as $1/A^{3/2}$ in the cold eras ($p=\tilde{p}=0$)

Therefore, in the radiative eras the source terms vanish and the evolution of the universe is simply driven by its self interaction. In the cold eras the source terms varies as $(A^{1/2} - A^{-1/2})\rho_0$ where ρ_0 is presumably the common positive density the two conjugate sides started from.

Thus we get simple evolution laws and solutions in the particular ranges for the scale factor ($A=a^2$): $a \ll 1, a \gg 1$ since the differential equations read:

$$a \ll 1 \Rightarrow (\ddot{1/a}) = \frac{-n\pi G\rho_0}{6} \Rightarrow a \propto \frac{-12}{n\pi G\rho_0} \frac{1}{t^2} \text{ where } t < 0, \quad (9)$$

$$a \gg 1 \Rightarrow \ddot{a} = \frac{-n\pi G\rho_0}{6} \Rightarrow a \propto \frac{-n\pi G\rho_0}{12} t^2 \text{ where } t > 0. \quad (10)$$

where we anticipate that the constant $n\pi G$ is negative.

In the radiative era, the trivial $A=1$ stationary solution describes a self conjugate world. A perturbation to this stationary solution, probably occurred for the birth of times to take place and see a couple of conjugate universes start evolving from $A=1, t=0$. These worlds have simple evolution laws in the particular ranges $A \ll 1, A \approx 1, A \gg 1$. Indeed, the scale factor evolution is then driven (here a non dimensional time unit is used) by the following differential equations in the three particular domains:

$$A \ll 1 \Rightarrow \ddot{A} = \frac{3}{2} \frac{\dot{A}^2}{A} \Rightarrow A \propto 1/t^2 \Rightarrow a \propto 1/t \text{ where } t < 0, \quad (11)$$

$$A \approx 1 \Rightarrow \ddot{A} = \frac{\dot{A}^2}{A} \Rightarrow A \propto e^t \Rightarrow a \propto e^{t/2}, \quad (12)$$

$$A \gg 1 \Rightarrow \ddot{A} = \frac{1}{2} \frac{\dot{A}^2}{A} \Rightarrow A \propto t^2 \text{ where } t > 0 \Rightarrow a \propto t. \quad (13)$$

Of course we can check that $t \rightarrow -t$ implies $1/t^2 \rightarrow t^2$ but also $e^t \rightarrow e^{-t}$ thus $A \rightarrow 1/A, B \rightarrow 1/B$ when t reverses for all our solutions Remarkably, we have $A^{-1}(-t) = A(t)$. The conjugate background is therefore obtained by time reversal. Two opposite time parameters joined each over at the conjugate sides $t=0$. But here time reversal does not mean going backward in time anymore. As shown in the first picture, reversing time twice can never make you reappear in the past in a given side of the universe. Not only our universe can be accelerated thanks to the $1/t^2$ solution (which translates into a t^2 evolution in standard comoving coordinates) without any need for a cosmological constant or dark energy component but it is flat without inflation and gets rid of the big-bang singularity at $t=0$. The $1/t$ and t radiative eras solutions respectively translate into e^t and $t^{1/2}$ evolutions in standard comoving coordinates.

We can also notice that the coordinate system where $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$ defines two fundamental scales or constants of nature. The first one is the speed of light $c=1$ in this system. On the second one $a(t=0) = 1$ depends the time of the transition between exponential regime and later regime. Considered as fundamental constants both are for the time being arbitrary but measurable. There is no

suppression effect of the background here as in GR in the vicinity of a local potential. But we shall show in a forthcoming section why atoms and planetary systems typical sizes and periods are affected in the same way so that it is not possible to locally detect any effect of the cosmological background on planet trajectories from our reference rods and clocks point of view. Only the comparison of atomic periods with free photons periods, i.e. the redshift, can allow us to probe the cosmological expansion.

A striking and very uncommon feature is that the evolution of the scale factor is driven by the gravitational interaction between the conjugated sides of the universe corresponding to the two forms independently of their matter and radiation content in the radiative era. Also in the cold era, probing the flatness or the variations of the expansion rate of our universe should no longer allow us to draw any conclusion about its content except in case among the three solutions, one side does not choose a single one everywhere, but instead is divided into many domains with different regimes of $A(t)$ depending on local density perturbations. This discussion will be reopened in a next section.

14. B=-A Perturbations: Gravitational Waves

The propertimes we can build with this field a priori take the form:

$$d\tau^2 = A(r, t) [d\sigma^2 - dt^2]$$

$$d\tilde{\tau}^2 = A^{-1}(r, t) [d\sigma^2 - dt^2]$$

The expressions will be simpler if we adopt the dynamical variable $h(r, t)$ defined by $A = e^{h(r, t)}$. In vacuum we obtain

$$\Rightarrow \square h(r, t) = \frac{\text{Tanh}(h)}{2} (h'^2 - \dot{h}^2)$$

having massless plane wave solutions.

$$h(r, t) = \sin(Et - \vec{p}\vec{r}); |\vec{p}| = E$$

The nonlinear term on the right hand side of the $h(r, t)$ dynamical equation vanishes for any propagating $h(r, t)$ or $h(r+t)$ but not for a superposition of such functions. For instance, the superposition of an outgoing (retarded) and ingoing (advanced) spherical waves with the same frequencies is a standing wave producing a nonzero nonlinear term which can act as a source term. Indeed, though on large scales the averaged $h'^2 - \dot{h}^2$ vanishes, on small space-time scales relative to the wavelength, the Zitterbewegung of $h'^2 - \dot{h}^2$ may have been the perturbation needed to start a non stationary evolution of the background, i.e. for the birth of a couple of time reversal conjugate universes which scale factors evolved as shown in the previous section. Note that such nonlinear effect are also efficiently reduced by the $\text{Tanh}(h)$ factor for small h . The basic small perturbation solution is thus a plane wave

$$A(r, t) = e^{\sin(Et - \vec{p}\vec{r})} \quad |\vec{p}| = E$$

We remind the reader that the three solutions for the cosmological $B(t)=-A(t)$ co-exist, so we have two evolving conjugate worlds as well as a third globally static one in between them. Independent solutions of the $B(r,t)$ field that we are studying now and which is itself two-sided can be produced and run through all three universes. Thus the conjugate wave

$$\tilde{A}(-r, -t) = e^{sin(-Et + \vec{p}\vec{r})}$$

has a negative frequency but is still propagating on the same side of the $B=-A$ cosmological field as $A(r,t)$. It also has a reversed momentum, so that the discrete symmetry involved here is space and time reversal (r,t to $-r,-t$). The rest energy reversal we got in the $B=-1/A$ gravity theory actually also went with time reversal of the $B=-A$ background field. So the discrete symmetry was also (r,t to $-r,-t$).

We then have two possibilities to interpret the negative frequency conjugate wave: The first possibility to consider is to deny any physical role to be played by this conjugate form because firstly it is not another degree of freedom but just the other side of the same object, secondly if it were the negative energy side of the field any interaction with (coupling to) a positive energy side of another field in the same universe would be forbidden and finally it is not propagating in the other universes which means that the positive energy wave propagating on our side is not at the same time propagating in the form of a negative energy wave on the conjugate side of the universe.

Another tempting interpretation is to consider that the conjugate form still being on our side must always be taken at $-t$ instead of t to compensate for the negative sign of its energy. In this case, it may be interpreted as a negative energy going backward in time equivalent to a positive energy antiwave (antiparticle) going forward in time, and we can even allow this object to couple to the positive energy form of other waves there.

We now demonstrate that these waves can be responsible for the decay of the binary pulsar. We follow Weinberg's computation of the power emitted per unit solid angle and adopt the same notations to obtain the same energy lost through our gravitational waves radiation by the binary pulsar as in GR in good agreement with the observed decay of the orbit period. Our understanding is that any distribution of matter can source the $B=-1/A$ field as we explained in previous sections, gravitationally on our side and antigravitationally on the conjugate side, but can also source in the three universes, independent $h(r,t)$ waves. In anyone of these universes their Lagrangian satisfies:

$$-n\pi GL = \sqrt{g}R + \frac{1}{\sqrt{g}}R_{g \rightarrow 1/g} = 3Cosh(h)(h'^2 - \dot{h}^2) + 6(Sinh(h))(\Delta h - \ddot{h})$$

The conserved Noether current being

$$t_{\mu\nu} = \frac{\partial L}{\partial(\partial^\mu h)}\partial_\nu h - \eta_{\mu\nu}L - \partial^\rho\left(\frac{\partial L}{\partial(\partial^\mu\partial^\rho h)}\right)\partial_\nu h + \frac{\partial L}{\partial(\partial^\mu\partial^\rho h)}\partial_\nu\partial^\rho h$$

we obtain

$$t_{00} = \frac{1}{-n\pi G} (3Cosh(h)(h'^2 - \dot{h}^2) + 6Sinh(h)\Delta h)$$

Thus for any propagating solution $h(r,t)$ or $h(r+t)$:

$$t_{00} = \frac{1}{-n\pi G} (6Sinh(h)\Delta h \approx 6h\Delta h = 6h\ddot{h})$$

the approximation being for weak $h(r,t)$ fields.

The radial momentum component of our gravitational wave energy momentum tensor reads:

$$t_{r0} = \frac{6}{-n\pi G} Cosh(h)h'\dot{h} \approx \frac{6}{-n\pi G} h'\dot{h}$$

We have used the genuine energy-momentum tensor derived from the Noether theorem which is not the pseudo-tensor of GR. It exploits the global Lorentz invariance of our flat spacetime theory. The energy carried by the field $h(r,t)$ is positive (the negative $n=-48$ as we shall show).

We now compute the energy radiated by the small $h(r,t)$ field in our $B(t)=-A(t)$ universe by the binary pulsar. The field is solution of $6(\Delta - \partial_0^2)h(r,t) = n\pi G\delta(r)\delta(t)$ when excited by a dirac impulse $\delta(r)\delta(t)$. Again, as for the $B=-1/A$ case we want to determine n that insures the compatibility condition in the static case: $h(r)=-2U(r)$ where $U(r)$ would again be solution of $\Delta U(r) = 4\pi G\delta(r)$. Thus now n equals -48 and our Lagrangian for gravity in the $B=-A$ sector is

$$L = \frac{1}{48\pi G} (\sqrt{g}R + \frac{1}{\sqrt{g}}R_{g \rightarrow 1/g})$$

For any extended non relativistic source δT_μ^μ (remember that our single equation only involved tensor traces) the solution is the "retarded potential":

$$h(x,t) = 2G \int d^3x' \frac{\delta T_\mu^\mu(x', t - |x - x'|)}{|x - x'|}$$

Replacing by the expression of our wave solution,

$$h = \sum_{\omega, k} h(\omega, k) e^{i(\omega t - kr)} + h^*(\omega, k) e^{-i(\omega t - kr)}$$

$$\langle t_{r0}(\omega, k) \rangle = \frac{1}{8\pi G} \langle h'\dot{h} \rangle_{\omega, k} = \frac{1}{4\pi G} \omega^2 |h(\omega, k)|^2$$

Allowing to get the total power emitted per unit solid angle in the direction of k :

$$\frac{dP}{d\Omega}(\omega, k) = r^2 \langle t_{r0}(\omega, k) \rangle = r^2 \frac{1}{4\pi G} \omega^2 |h(\omega, k)|^2 = \frac{G}{\pi} \omega^2 |\delta T_\mu^\mu|^2(\omega, k)$$

Then following Weinberg, we use $T_{0i}(k, \omega) = -\hat{k}^j T_{ji}(k, \omega)$ and $T_{00}(k, \omega) = \hat{k}^i \hat{k}^j T_{ji}(k, \omega)$ (where $\hat{k}$ is the unit vector in direction of the vector k), assume that the source radius is much smaller than the wavelength $1/\omega$ to write $T_{ij}(k, \omega) \approx -\frac{\omega^2}{2} D_{ij}(\omega)$ to obtain in terms of the moment of inertia Fourier transforms $D_{ij}(\omega)$:

$$\frac{dP}{d\Omega}(\omega, k) = \frac{G\omega^6}{4\pi} (\hat{k}_i \hat{k}_j \hat{k}_l \hat{k}_m + \delta_{ij} \delta_{lm} - 2\hat{k}_i \hat{k}_j \delta_{lm}) D_{ij}^*(\omega) D_{lm}(\omega)$$

The next step is to do the integral over solid angle

$$P = \left(\frac{1}{15} (\delta_{ij}\delta_{lm} + \delta_{il}\delta_{jm} + \delta_{im}\delta_{jl}) + \frac{1}{3} \delta_{lm}\delta_{ij} \right) G\omega^6 D_{ij}^*(\omega) D_{lm}(\omega)$$

$\delta_{ij} D_{ij}$ terms do not contribute for a rotating body and we get

$$P = \frac{1}{15} G\omega^6 [2D_{ij}^*(\omega) D_{ij}(\omega)] = \frac{1}{15} G\omega^6 8D_{11}^2(\omega)$$

For a rotating body with angular speed Ω , equatorial ellipticity e , moment of inertia I in the rotating coordinates, $\omega = 2\Omega$, $D_{11}(\omega) = \frac{eI}{4}$ and the radiated power reads:

$$P = \frac{8}{15} 64G\Omega^6 e^2 \frac{I^2}{16} = \frac{32}{15} G\Omega^6 e^2 I^2$$

We find that an extra factor three eventually is needed to get the same lost total power $3P$ as in General Relativity. We interpret this as a strong indication that the h fields exists in 3 kinds. Though it is not yet already obvious which kind of symmetry is involved here, the analogy with the three mesons π^+ , π^- , π^0 , is attractive. These are Lorentz scalars just as the field Φ we may have defined by $g_{\mu\nu} = \Phi\eta_{\mu\nu}$ and even though they are massive while DG fields are massless, we shall explain later how our fields are expected to acquire mass.

The main difference as opposed to the GR case is that our gravitational wave is found to propagate pure scalar modes: $-g_{00} = g_{11} = g_{22} = g_{33} = e^h$. The wave clearly affects all spatial directions in the same way so that no interferometer will be able to detect it. To see this we can for instance adopt the experimenter proper time as our new time coordinate t' and in this new coordinate system the only non vanishing field perturbation elements are $h_{11} = h_{22} = h_{33}$ depending on t' and the direction of propagation z . The Riemann tensor elements we need to characterize the observational effects of our wave are R_{0i0j} with only non vanishing elements in the small h approximation appropriate for a wave, $R_{0i0i} = \frac{1}{2}\ddot{h}_{ii}$. Hence we have the same effect on any arm of an interferometer whatever the propagation direction of the wave.

The h field once quantized is expected to generate a new gravitational propagated interaction in addition to the Schwarzschild non propagated solution we obtained in the previous section. We cannot add its potential to the Schwarzschild one since this would severely conflict with observations except if the B=A solution is quantized with another Planck constant much larger than the one used to quantize electromagnetic interactions. Indeed, it was pointed out by J.M Souriau that the planetary periods in the solar system are Fibonacci multiples of an approximate 30 days period (see Ref. 48), which we interpret in the framework of our quantum gravity with huge gravity quanta exchanges (see Ref. 49 and 50 with references therein). Postulating an energy cutoff for the virtual gravitons (DG itself is a suitable framework to generate such a cutoff as we explain in a next section) at distances smaller than a given threshold the force becomes negligible. Following this way of thinking an extra contribution to gravity might only arise beyond at least interstellar distances (this was also postulated in Ref. 98) since no deviation

from Newtonian gravity was observed at smaller distances. At smaller distances the extra interaction would thus exhibit a kind of asymptotic freedom. Notice that the cutoff only applies to virtual gravitons, not the real ones emitted in the binary pulsar gravitationnal radiation.

An even simpler proposal becomes possible in case the fields are understood to acquire mass and we may even identify our field to be that of pion particles. Indeed, in this case the Yukawa potential e^{-mr}/r makes the interaction a very short distance one and its potential completely negligible as compared to the nuclear interactions taking place at the same scales. On the other hand, adding a mass term is possible without introducing any new dynamical degree of freedom as is unavoidable when one adds massive gravitons in GR so our result for the power emitted through gravitational waves radiation should not change. Following this way of thinking it is also tempting to consider the validity of our wave equations at the nuclear scale with another coupling constant to account for the Yukawa interaction though nowadays the latter is rather understood to be an effective description, QCD being the accepted fundamental theory for the strong interactions.

15. Pioneer Effect and Pseudo-Horizon Effect

15.1. *The Pioneer Effect*

The so called Pioneer anomaly arose as an anomalous drift in time of the radiowave frequency received from both Pioneer 10 and 11 as compared to reference clocks on earth (Ref. 17). The anomalous drift was found constant over more than 10 years and from 40 to 60 AU

$$\frac{\dot{f}}{f} = (5.6 \pm 0.9)10^{-18}/s$$

including in the error all systematical sources of uncertainties. Since all possible systematical origins investigated by Anderson and collaborators (Ref. 17) were excluded or found very unlikely to account for the Pioneer anomaly, there is increasing evidence that it is now of fundamental physics origin. In GR, the Pioneer anomaly is a fatal one. Indeed, not only the sign of the shift is opposite to what we would have expected from a background effect but moreover the best effort in GR to melt together a background metric with a Schwarzschild metric is the McVittie metric (see Ref. 42 and references therein):

$$d\tau^2 = -\frac{1 - \frac{Gm}{2ra(t)}}{1 + \frac{Gm}{2ra(t)}} dt^2 + \left(1 + \frac{Gm}{2ra(t)}\right)^4 a^2(t) d\sigma^2 \quad (14)$$

where $a(t)$ can be absorbed by a coordinate transformation $ra(t) \rightarrow r$ so that the background is completely suppressed except in the space-time g_{ti} elements where its effect is at a much lower level than in the reported anomaly magnitude. In the combined DG solution, there is superposition without suppression:

$$d\tau^2 = -e^{-\frac{2Gm}{r}} a^2(t) dt^2 + e^{\frac{2Gm}{r}} a^2(t) d\sigma^2 \quad (15)$$

While in such $A(t)=-B(t)$ background, photons keep unaffected, the reference atomic periods contract or expand (but $B(t) = -A(t)$ also appears to rescale masses (Ref 42)) resulting in an apparent cosmological red or blue shift. Therefore the Pioneer photons periods should also appear shifted with respect to atomic clock references due to the $B=-A$ background but due to their very small time of flight, the effect is very tiny. At best it would result in the good sign uniformly changing shift but with a rate too low by a factor $\frac{v}{c}$ to account for the reported anomaly showing up from the Pioneer spacecrafts data. However we are well motivated to keep searching for a Background effect solution because of the strange coincidence $\frac{\dot{f}}{f} = 2 \frac{1}{11.10^9 \text{years}} \sim 2 \frac{\dot{a}}{a} \equiv 2H_0$ at a roughly 15 percent confidence level coming from both the uncertainty in the directly measured Hubble parameter H_0 and in the effect.

Several authors have argued that, because of its weakness, the Pioneer anomaly seen as such a time dependent metric effect (due to the lacking of any fundamental theory to account for such a form the issue is always examined from a phenomenological point of view), a clock acceleration rather than a mere acceleration, would actually entail no contradiction with the four classical tests of General Relativity since the trajectories are unperturbed (Ref. 45, 46 and references therein). At the contrary, all theories appealing to extra accelerations to produce a Doppler effect in order to explain the Pioneer anomaly are in serious conflict with recent tests of the outest planets trajectories in the solar system. An exception may be a modification of the r dependence of space-space metric elements having a very little influence on almost circular planet trajectories but accelerating significantly radial Pioneer like trajectories as proposed in Ref. 83 (see also Ref. 84). But this is an ad-hoc fine-tuned model.

For significant effects to appear in DG we are led to consider the case where the emitter and receiver are not in the same background which happens if somewhere between the earth and the spacecraft there is a discontinuity of the background. Indeed the received radiowave signal frequencies are electronically shifted before reemission by the spacecraft antenna in such a way that these should undergo an $a^2(t)$ drift as compared to earth standards if at a discontinuity between the earth and Pioneer spacecraft, the $B = -A$ background field has flipped according $A(t) \rightarrow 1/A(t)$ producing an $a^2(t)$ Pioneer clocks drift with respect to earth clocks on the other side of the discontinuity. To show this, we can write down the following metrics on earth and on Pioneer to get the clocks drift in time relative to each other. On Pioneer:

$$-d\tau^2 = a^2(t)(dt^2 - d\sigma^2) \Rightarrow dt_{\text{Pioneer clock}} \propto \frac{1}{a(t)}d\tau \quad (16)$$

On Earth:

$$-d\tau^2 = a^{-2}(t)(dt^2 - d\sigma^2) \Rightarrow dt_{\text{Earth clock}} \propto a(t)d\tau \quad (17)$$

The result is a frequency acceleration of Pioneer clocks relative to our earth clocks $f_{\text{Pioneer}} = a^2(t)f_{\text{earth}}$ yielding the anomalous $\frac{\dot{f}_{\text{Pioneer}}}{f_{\text{Pioneer}}} = 2H_0$. Therefore,

there must be a discrete jump from $a(t)$ to $1/a(t)$ of the background field between us and Pioneer so that the effect could only start to be seen after the crossing of this frontier by the spacecrafts. Within the error bars the jump (see the steep rise up of the effect in Ref. 17) could not have been better evidenced than it was around 15 AU in 1983 by Pioneer 11.

The Pioneer anomaly in our framework is obviously an LPI in time violation (thus an SEP violating effect), so the simple superposition of the background and perturbation readable in DG's solution might be in conflict with the usual tests of LPI in time expressed as a variation of the gravitational constants G which relative variations are now constrained at a level much lower than H_0 . These tests were performed in the inner part of the solar system. We shall reconsider this issue later after a careful study of the behaviour of an atom in the DG background field to prove that if planet trajectories and our local reference atomic rods and clocks are in the same background there is no detectable effect at all, and therefore no conflict with constraints on the time variation of G . We are mainly concerned here with the effect on the outer solar system trajectories with respect to our local clocks which are not in the same background. We find that the apparant effect of reduction of the outer planet period is too small to be detected since the outer planet trajectories were not monitored by accurate Doppler or Laser Ranging methods but simply by following their visual position in the sky. By far, the best precision on the pericenter precession according Ref. 73 were a 0.036 Arcsec/century for Jupiter to be compared with the 0.0024 Arcsec/(10years) expected in DG from 1997 to 2007 if, as we believe, the discontinuity was between the earth and Jupiter during that period (planet and our earth clocks not in the same background).

Photons are not sensitive to an homogeneous background $B=-A$ field except at a discontinuity which may allow them to jump to the conjugate side. For matter, the discontinuity represents a potential barrier that might produce caustic like effects where matter is trapped along the discontinuity forming matter rings or shells. The seemingly time dependent flux of Ultra High Energy cosmic rays as a consequence of the discontinuity drift in time in our vicinity is also an interesting possibility if some of these particles originate somehow at the gravitational discontinuity.

The effect on local rods might reverse periodically as we shall show, so that integrated along a cosmological path, its impact is not trivial. Thus distant photons periods evolution relative to local rods may alternate between a t^2 regime and $1/t^2$ regime. The amount of time spent in the two different regimes will determine the long term behaviour and the expected Hubble diagram parameters.

15.2. *The Pseudo Black Hole Horizon*

Our exponential solution tells us that there is no more BH horizon in DG. However a test mass approaching the Schwarzschild radius of a massive body is propagating

in the background and perturbation superposition:

$$-d\tau^2 = \frac{1}{A(r)}a(t)^2 dt^2 - A(r)a(t)^2 d\sigma^2 \quad (18)$$

But for a gravitational field such that the $A(r)$ term "crosses" the background term we locally get $g_{00} = -1$ and a lowest achievable potential. For a test mass at rest ($d\sigma = 0$) reaching this radius, the "feeled metric" is $-d\tau^2 = dt^2 = dt^2 - d\sigma^2$ and we postulate that this is a necessary and sufficient condition to allow a local spontaneous phase rotation of the gravitational field transferring the test mass in the Euclidian world and possibly via an extra rotation in the conjugate universe. As well as a genuine black hole horizon this mechanism would account for the absence of thermonuclear bursts when matter falls down into BH candidates.

However, depending on $A(r)$ and $a(t)$, it is possible that no particular r_0 will eventually allow these terms to cross each other. Instead, we can have $A^{-1}(r_0)A(t) = 1$ but then the crossing can only happen and the primordial symmetry be recovered provided the field jumps from $A(t)$ to $1/A(t)$ at r_0 . Such kind of discontinuity is believed to be at the origin of the Pioneer anomaly as explained in the previous section. Interestingly, r_0 should propagate to maintain:

$$e^{Gm/r_0(t)} = a(t) \quad (19)$$

Over a negligible timescale compared to the universe age and in the weak field approximation, the propagation time $t_2 - t_1$ from r_1 to r_2 is given by:

$$H_0(t_2 - t_1) = \phi(r_2) - \phi(r_1) \quad (20)$$

where $\phi(r)$ stands for the gravitational potential at r . We find that the discontinuity takes 26000 and 9 years to travel through the solar and earth potential respectively e.g from vanishing potential at infinity to the potential reached at the surface of the objects. The Pioneer effect seems to tell us that the discontinuity was at around 12.5 A.U from the sun in 1983, the time when Pioneer 11 detected the very steep rise up of the effect. It follows that it must have reached the Jovian potential well and started to fall inside it in 1997 while it will reach the earth potential in 2104. It is thus now already falling in the potential wells of Jupiter and the outer planets Saturn, Uranus and Neptune, which might be correlated with the unexplained activity of Saturn's satellite Encelade and perturbed magnetic field of Uranus.

Recently a Bubbling behaviour accompanied by ripples visible in the hot gas of the Perseus cluster has been reported by Chandra Ref. 25. We interpret this as the first observation of a periodic generation of gravitational discontinuities which in turn produce our scalar (under rotation in a plane perpendicular to the direction of propagation) longitudinal gravitational wave ripples rather than sound waves as interpreted in Ref. 25. The Bubbling frequency is also the gravitational waves frequency which can be computed accurately knowing the measured wavelength (approximately 11kpc) and the GWs speed which is the speed of light. One obtains that the frequency is just 2^{49} times lower than 475 cycles per second, the frequency

corresponding to B-flat above middle C and the corresponding period is roughly 30000 years which compares very well with the time needed by the discontinuity to travel across the gravitational potential (from infinity to the surface of the star) of a typical star as is our sun. Therefore 30000 years is a good estimation of the time needed for a gravitational discontinuity travelling through the almost flat potential of a galaxy to scan the gravitational potential of the majority of stars in the galaxy (only a minority of white dwarf stars have much larger surface potentials). This seems to indicate that when most of the stars surfaces have been reached by the discontinuity, an instability makes the discontinuity jump back or bounce somewhere in the deeper potential near a galaxy center to initiate another 30000 years path.

When the discontinuity reaches the vicinity of a planet or star, the isopotential is roughly a spherical surface surrounding the object. An isopotential where we find a discontinuity or a crossing of the two forms of the metric has an other interesting effect: because it acts as a boundary for a spatial domain, the asymptotic decrease of local field perturbations at infinity required by the Gauss theorem is a condition which may not be possible to fulfill anymore, so that the finite field value at the boundary might behave as a central effective source mass. This mechanism might create the illusion of a dark several billion solar masses object with huge effects on stellar dynamics in the central arcsecond of our galaxy as reported in Ref. 47.

16. General equations of motion: toward the unification of gravity and electromagnetism

With the metric in our usual isotropic form, the free fall equations of motion read

$$\begin{aligned} \frac{d^2 r}{dp^2} + \frac{1}{2} \frac{A'}{A} \left(\frac{dr}{dp} \right)^2 - \left(\frac{d\theta}{dp} \right)^2 \left(\frac{A'}{2A} r^2 + r \right) - \left(\frac{d\phi}{dp} \right)^2 \sin^2 \theta \left(\frac{A'}{2A} r^2 + r \right) + \frac{1}{2} \frac{B'}{A} \left(\frac{dt}{dp} \right)^2 + \frac{\dot{A}}{A} \frac{dr}{dp} \frac{dt}{dp} &= 0 \\ \frac{d^2 \phi}{dp^2} + \frac{\dot{A}}{A} \frac{d\phi}{dp} \frac{dt}{dp} + 2 \left(\frac{A'}{2A} + \frac{1}{r} \right) \frac{dr}{dp} \frac{d\phi}{dp} + 2 \cot \theta \frac{d\theta}{dp} \frac{d\phi}{dp} &= 0 \\ \frac{d^2 \theta}{dp^2} + 2 \left(\frac{A'}{2A} + \frac{1}{r} \right) \frac{dr}{dp} \frac{d\theta}{dp} - \sin \theta \cos \theta \left(\frac{d\phi}{dp} \right)^2 + \frac{\dot{A}}{A} \frac{d\theta}{dp} \frac{dt}{dp} &= 0 \\ \frac{d^2 t}{dp^2} + \frac{B'}{B} \left(\frac{dr}{dp} \right) \left(\frac{dt}{dp} \right) + \frac{1}{2} \frac{\dot{B}}{B} \left(\frac{dt}{dp} \right)^2 + \frac{1}{2} \left(\frac{d\phi}{dp} \right)^2 \frac{\dot{A}}{B} r^2 \sin^2 \theta + \frac{1}{2} \left(\frac{dr}{dp} \right)^2 \frac{\dot{A}}{B} + \frac{1}{2} \left(\frac{d\theta}{dp} \right)^2 \frac{\dot{A}}{B} r^2 &= 0 \end{aligned}$$

With $d\theta = 0$, $\theta = \pi/2$, θ is not anymore a dynamical variable, expliciting the spatial and temporal parts : $A(r,t) = A(r)a^2(t)$, $B(r,t) = B(r)a^2(t)$, where $B(r) = 1/A(r)$ stands for our time independent Schwarzschild solution.

$$\begin{aligned} \frac{d^2 r}{dp^2} + \frac{1}{2} \frac{A'}{A} \left(\frac{dr}{dp} \right)^2 - \left(\frac{d\phi}{dp} \right)^2 \left(\frac{A'}{2A} r^2 + r \right) + \frac{1}{2} \frac{B'}{A} \left(\frac{dt}{dp} \right)^2 + 2 \frac{\dot{A}}{a} \frac{dr}{dp} \frac{dt}{dp} &= 0 \\ \frac{d^2 \phi}{dp^2} + 2 \frac{\dot{A}}{a} \frac{d\phi}{dp} \frac{dt}{dp} + 2 \left(\frac{A'}{2A} + \frac{1}{r} \right) \frac{dr}{dp} \frac{d\phi}{dp} &= 0 \\ \frac{d^2 t}{dp^2} + \frac{B'}{B} \left(\frac{dr}{dp} \right) \left(\frac{dt}{dp} \right) + \frac{\dot{A}}{a} \left(\frac{dt}{dp} \right)^2 + \left(\frac{d\phi}{dp} \right)^2 \frac{A}{B} \frac{\dot{a}}{a^2} r^2 + \left(\frac{dr}{dp} \right)^2 \frac{A}{B} \frac{\dot{a}}{a^2} &= 0 \end{aligned}$$

In our privileged frame, light propagation is obviously not sensitive to the scale factor evolution. In the following we are mainly interested in the $a(t)$ leading terms influencing typical sizes and times of non relativistic bound systems. So we work

with slow velocity approximated equations. We integrate the second equation after dividing it by $d\phi/dp$

$$\frac{d}{dp} \left[\ln \left(\frac{d\phi}{dp} \right) + \ln A + \ln a^2 + 2 \ln r \right] = 0 \Rightarrow \frac{d\phi}{dp} \propto \frac{1}{Aa^2r^2}$$

In the small speed approximation we can neglect the $(a(t)dr/dt)^2$ terms and after dividing by dt/dp the third equation integration leads to

$$\frac{d}{dp} \left(\ln \frac{dt}{dp} + \ln B + \ln a \right) = 0 \Rightarrow \frac{dt}{dp} \propto \frac{1}{aB}$$

The first equation in the small speed limit and using $\frac{d}{dt} \left(\frac{dt}{dp} \right) = -\frac{\dot{a}}{a} \frac{dt}{dp}$ gives

$$\begin{aligned} \frac{d}{dt} \left[\frac{dt}{dp} \frac{dr}{dt} \right] \frac{dt}{dp} + \frac{1}{2} \frac{B'}{A} \left(\frac{dt}{dp} \right)^2 + 2 \frac{\dot{a}}{a} \dot{r} \left(\frac{dt}{dp} \right)^2 &= 0 \\ \Rightarrow \left[\ddot{r} - \frac{\dot{a}}{a} \dot{r} \right] + \frac{1}{2} \frac{B'}{A} + 2 \frac{\dot{a}}{a} \dot{r} &= 0 \end{aligned}$$

Therefore

$$\ddot{r} = -\frac{1}{2} \frac{B'}{A} - \frac{\dot{a}}{a} \dot{r} \approx -\frac{Gm}{r^2} - \frac{\dot{a}}{a} \dot{r}$$

Using $\frac{dt}{dp} = \frac{1}{aB}$ to eliminate dp also gives $\dot{\phi} = \frac{B}{Aa^2r^2}$

The effect of the expansion on light or gravitationally bound system or any hyperbolic trajectory can of course only be probed with respect to the effect of expansion on an electromagnetically bound system: the reference atomic sizes and periods.

The equations of motion of any body of charge q and mass m in a gravitational field and electric potential field $V(r, t)$ are the one written above which already include in the gravitational field the effect of a local static gravitational potential and the scale factor evolution and must now be corrected for an extra electric acceleration $[d^2x^\mu/dp^2]_{EM} = \frac{e}{m} g^{\mu\rho} (\partial_\rho A_\nu - \partial_\nu A_\rho) \frac{dx^\nu}{dp}$ where $A_\mu = (V(r, t), 0, 0, 0)$. Explicitly

$$\begin{aligned} \left[\frac{d^2r}{dp^2} \right]_{EM} &= \frac{q}{m} \frac{V(r, t)'}{Aa^2(t)} \frac{dt}{dp} \\ \left[\frac{d^2\phi}{dp^2} \right]_{EM} &= 0 \\ \left[\frac{d^2t}{dp^2} \right]_{EM} &= \frac{q}{m} \frac{V(r, t)'}{Ba^2(t)} \frac{dr}{dp} \end{aligned}$$

The $d\phi/dp$ equation is thus integrated as in the purely gravitationnal case to give the same result. We now take it for granted that $V(r, t)$ on its own side also includes the effect of expansion and shall show later that $V(r, t) = a(t)V(r)$ where $V(r)$ stands for the usual electrostatic potential.

We first perform the integration of

$$\frac{dt}{dp} \frac{d}{dp} \left(\ln \frac{dt}{dp} + \ln B + \ln a \right) = \frac{q}{m} \frac{dV}{B(r)a(t)dp} \quad (21)$$

or defining $u = B(r)a(t)\frac{dt}{dp}$:

$$u \frac{d}{dp} (\ln u) = \frac{q}{m} \frac{dV}{dp} \quad (22)$$

hence now $dt/dp = \frac{1}{a(t)B(r)}(\frac{q}{m}V(r) + 1)$ and in the case of atomic energy levels $dt/dp \approx \frac{1}{a(t)B(r)}$. Replacing as before the first equation simplifies and eventually we have the two equations of motion including all effects :

$$\begin{aligned} \ddot{r} &= \frac{q}{m}V'(r)\frac{B}{A}(\frac{q}{m}V(r) + 1) - \frac{1}{2}\frac{B'}{A} - \frac{\dot{a}}{a}\dot{r} \approx \frac{q}{m}V'(r) - \frac{1}{2}\frac{B'}{A} - \frac{\dot{a}}{a}\dot{r} \\ \dot{\phi} &= \frac{B}{Aar^2}(\frac{q}{m}V(r) + 1) \approx \frac{B}{Aar^2} \end{aligned}$$

where the approximative results are given in the case of atomic energy levels.

We can now interpret these results.

- The time evolution of ϕ for an atom tells us that the characteristic period of a clock behaves as $a(t)$. Let us write this $dt_{clock} \propto a(t)dt$. Since $d\tau \propto a(t)dt$, the time given by physical clocks is just the proptime.
- Analysing the period of a distant Supernova radiation with respect to such a clock, it is straightforward to check that in this time coordinate, it appears redshifted by an accelerated t_{clock}^2 (resp decelerated $t_{clock}^{2/3}$) scale factor in case $a(t) = 1/t^2$ (resp $a(t) = t^2$). See Figure 2. This result is most easily obtained by a change of variable from the preferred time t to t_{clock} . Under this transformation the expression of the background field metric which was

$$-d\tau^2 = (1/t^2)^2(dt^2 - d\sigma^2) \quad (23)$$

becomes

$$-d\tau^2 = dt_{clock}^2 - t_{clock}^4 d\sigma^2 \quad (24)$$

so the cosmological phenomenology is just given by the scale factor $a(t_{clock}) \propto t_{clock}^2$ while $a(t_{clock}) \propto t_{clock}$ corresponds to the exponential regime of $a(t)$.

- We find that the $a(t)$ dependency of r and ϕ for a bound system do not actually depend on the nature of the bounding force but rather on the radial speed $\dot{r}$ only. This means that for circular orbits there is no way to see the effect of expansion by studying circular or small excentricity planet trajectories (with respect to our reference atom standards). Our assumption that $V(r, t) = a(t)V(r)$ was a crucial one to obtain this result. Had $V(r, t)$ been static, we would have obtained an $a(t)$ dependency in the final electric acceleration but not the gravitationnal one, a dead end since this would have led to effects usually referred to as variation of the constant G effects well constrained at the level of a percent of the Hubble factor (notice however that $\dot{\phi}$ behaves as in the standard picture detailed in Ref. 42, hence there is nothing really new as for the characteristic time

periods and energy levels). In the preferred coordinate system the atomic and gravitationally bound systems are only subjected to the tiny effect of $-\frac{\dot{a}}{a}\dot{r}$ relative to the unaffected light wavelengths. In the physical observer clocks coordinate system all are submitted to the same expansion effects except for the tiny friction effect.

- The $H_0\dot{r}$ acceleration must be compared with H_0c needed to account for the Pioneer anomaly if it could be interpreted as an acceleration. It is roughly 3.10^4 smaller so that obviously it could not have been detected by solar system spacecrafts up to now. Because $\dot{r}$ vanishes on a full period no advance of perihely effect can be seen. But on the long term a variation of the excentricity might be searched for with foreseenable long term effects on our earth climate. On the short term, because planets and satellites radial speeds are so small in the solar system $H_0\dot{r}$ is certainly too small to be detected since it doesn't lead to an integrative effect.

We can now justify the crucial $V(r, t) = a(t)V(r)$ assumption. This result is obtained when one realises that in DG the equations of the B=-A field are not only the gravitationnal waves ones but could also almost well describe the generation of the electromagnetic field with the appropriate coupling constant of course. The logarithm of the B=-A field is a new field whose elements satisfy linear wave equations. For instance in the static case, the equation satisfied by $V=\ln(B)$ is just the Poisson equation. So a possible link between gravity and electromagnetism might be something like a perturbative developpment (we let this as an open issue for the time being). But now, let us consider this new B=-A field instead of the former. The total corresponding field has diagonal components $B=V(r)a^2(t)=-A$. It just remains to notice that $g_{\mu\nu} = A_\mu^\alpha A_\nu^\beta \eta_{\alpha\beta}$ in terms of a vielbein field A_μ^α and identify A_μ^0 with the electromagnetic potential entering in our above propagation equations in the privileged frame. From $A_\mu^0 = (V(r)a(t), 0, 0, 0)$ our earlier assumption is justified and we by the way realise that this simple road toward unification of gravity and electromagnetism made possible by the B=-A DG sector is necessary to avoid being in conflict with accurate tests constraining the variation of G...

17. Mass, Cosmological Constant Terms, Divergences

Since one of our main motivations for DG was to avoid an epicyclic construction, cosmological constant terms are a priori forbidden in our framework unless these are unavoidable from a fundamental particle physics point of view. Thus, the question arises wether we should expect this kind of contribution to our gravitational sources from either symmetry breaking or vacuum quantum fluctuations.

It was shown in Ref. 6 that a left-handed kinetic and interaction Lagrangian can satisfactorily describe all known physics except mass terms which anyway remain very problematic in modern physics. This strongly supports the idea that the right handed chiral fields are not on our side : the most straightforward way, as explained in Ref. 6 to explain maximal parity violation observed in the weak interaction.

According to this idea, there would be no fundamental massive fields and mass would always be effective as is the mass of a photon propagating in a transparent medium. In particular we would like to give up the idea that mass may be generated by a symmetry breaking mechanism based on the interaction with an extra field as is the Higgs field. Therefore we are well motivated for trying to investigate the possibility of a vacuum ether which would be responsible for generating the effective masses of the fundamental massless fields propagating through and interacting with this medium. Of course work is needed to build the postulated ether and new mass generating mechanisms but we will start to show in a forthcoming section why it is clearly easier to follow this line of thinking in our framework. Of course one then avoids vacuum energy contributions from spontaneous symmetry breaking.

Now let us investigate quantum fluctuations. One usually assumes that there is a cut off at the unification scale or at the Planck scale. According to Ref. 99, if care is taken to regularize the UV divergent integrals in a Lorentz invariant way, only massive fields (mass m) vacuum quantum fluctuations contribute to a source term in the Einstein equation and this is a cosmological constant term. Thus our main hope is to completely avoid such contributions if our fundamental fields are only massless as we explained earlier.

However may be could we live more easily with huge vacuum quantum fluctuation cosmological constant terms than we can in standard model cosmology. Indeed, one important result we got is that whatever its value the source term enters our cosmological scale factor solutions $a(t)$ in such a way that it only defines a particular time scale $T = \sqrt{\frac{-12}{n\pi G(\rho_{vac} - 3p_{vac})}}$, the threshold (t must be $\gg T$) for the validity of our $a(t) \ll 1$ and $a(t) \gg 1$ solutions. Thus there remains only one serious issue : is the vacuum fluctuations contribution to $\rho_{vac} - 3p_{vac}$ finite ? We could also rely on Planck scale physics to get finite vacuum fluctuations contribution but our framework also seems to come with its own interesting possibilities to be explored.

For instance, if the barrier between the two conjugate worlds can be jumped over by particles above a given energy threshold, then QED loop divergences hopefully might cancel thanks to the positive and negative energy virtual propagators compensation. Such reconnection might take place through allowing particles to jump from one form to the conjugate one presumably through discontinuities where the conjugate forms meet each other.

Alternatively, we already saw that the time reversed wave taken at $-t$, $\tilde{g}_{\mu\nu}(-t)$ may well describe an anti wave. But if we follow our first possible interpretation the wave has a negative energy and is on our side. We didn't allow it to couple to positive energy waves because it is a conjugate form but if for some reason (let us keep open minded), above a given energy threshold, the direct and conjugate forms flip and exchange their roles and if this can occur for any kind of wave, since propagators for any virtual interaction (electromagnetic, weak or strong) boson exchange are opposite for opposite energy virtual carriers, their UV contributions above threshold hence their UV divergences should cancel out leading to a finite

theory.

18. Phenomenological Outlooks: Cosmology

We now show that this Dark Gravity Theory is a very promising alternative to Dark Matter Models. Of course, if we expect discontinuities of the background to be everywhere present in the universe, the first task before doing cosmology should be a detailed study of the phenomenology associated with these discontinuities to be able to predict where and when do they occur in the universe, if they affect the element abundances and so on. In this section we will just briefly consider the main observables of cosmology trying to anticipate whether the new components involved in our formalism might help us to understand the universe and provide a promising alternative to the concordant model which being effective anyway should not be considered as more than a good parametrisation of our ignorance. If we are distrustful of the standard model construction, this attitude is even more supported by recent works (Ref. 80) calling into question the cosmological principle and showing that the background inhomogeneous evolution is completely unpredictable due to GR non linearities so that the associated back-reaction effects might mimic any of our effective parametric models (any mixture of Dark energy and Dark matter)! In the following subsections we follow the scenario of a succession of alternating t_{clock}^2 and $t_{clock}^{2/3}$ regimes in the cold era and a succession of alternating $t_{clock}^{1/2}$ and $e^{t_{clock}}$ regimes in the radiative eras. From now on we remain in the observer coordinate system whose time t_{clock} is the proptime.

18.1. The Hubble Test of the Accelerating Universe

The final answer from the Hubble test will obviously depend on our ability to separate possibly new evolution effects driven by the physics of discontinuities in Supernovae explosions. We obtain the correct Pioneer blue-drifting shift by assuming that the earth were recently in the t^2 regime while Pioneer spacecrafts were in the $1/t^2$ regime. Therefore the Pioneer anomaly tells us that we are presently undergoing a t^2 regime leading to the decelerating universe so this solution could not take place during most of the universe recent history according to the SN data that show acceleration at least from $z=0.6$ to the present time. Therefore we suspect an alternate regime of the two solutions (power 2 and 2/3) manifesting itself as a mean t_{clock}^α regime, α depending on the relative time spent in the two regimes on large time scales. Let's try to justify this scenario: Discontinuities are scanning all gravitational potentials in the universe. A possibility is that reaching an unknown threshold potential level makes all discontinuities everywhere simultaneously stop their centripete propagation, makes the background field jump back and reinitialize to its asymptotic and primordial state (η) thus the discontinuities jump back as well to the asymptotic potential (zero potential level) and initiate a new cycle. We are confident in this idea because sometimes lighting balls explode which means that the discontinuity that surrounds them sometimes disappears suddenly rather than

scans the object from the periphery to the center making it evaporate progressively. Also notice that this instantaneous jump will have no effect by itself on atoms (sizes and frequencies) anywhere apart from changing the background evolution regime they will now feel.

Another more involved possibility and not exclusive of the former is to have various discontinuities scanning local potentials (not total and global potential) with various such thresholds thus different cycle periods corresponding to local potential wells.

If the threshold is as deep as a black hole potential near the horizon the period of such cycle exceeds the age of our universe while if the threshold corresponds to the typical potential level at the surface of most stars, the period would be about 30000 years. In case of a global threshold, the typical maximum potentials encountered on the largest scales might determine the threshold level: the peculiar speeds of galaxy clusters can reach typical 1000km/s values, meaning that gravitationnal potentials at least about 10 times larger than our sun surface potential are at work at these scales, which translates into 300000 years for the completion of one such cycle.

Anyway, if we imagine for instance that a local discontinuity runs through our solar system with a 26000 years period, an atom on earth appears to be at the edge of the potential well of the sun, in the sense that the time needed by a discontinuity to scan the potential difference between that potential at the earth position and the one found at the sun surface is much larger (26000 years) than the time needed to scan the potential difference between the asymptotic value of the potential and the sun potential near earth (150 years). Therefore, over cosmological time scales this discontinuity is more than 99 percent of the time between the earth and the sun. But this does not imply that our reference earth clocks drifts are mainly determined by one of the two possible $B(t)=-A(t)$ field regimes while an atom in the vicinity of the sun surface would mainly be submitted to the other regime. Such scenario would produce huge shifts on cosmological time scales between frequencies of atoms far away from the stars and near the stars and very large and common redshift anomalies. Though there are claims of such redshift anomalies, these are very uncommon (for instance between quasars and galaxies that seem to be interacting though at incompatible redshifts).

The solution to avoid such not commonly seen anomalies is already in our hands: indeed after the discontinuity has jumped back to the asymptotic potential, the roles of the two regimes are exchanged upstream and downstream from the discontinuity as compared to the previous cycle. Alternatively, it is possible that discontinuities propagate in pairs with a small constant potential difference separating them as is also suggested by recent observations by New Horizons.

Then on the mean, all atoms will have experienced the same background shifts of their frequencies on time scales comparable to the age of the universe because they will have spent approximately the same amount of time in the two background regimes.

Of course, there should remain anomalous fluctuations most easily detectable

in regions where the cycle periods are longest. The expected fluctuations because all stars may not be at the same stage of their cycles evolution should produce a small dispersion in the redshifts, a new systematical but very small effect in the Hubble diagram most easy to detect with very low redshift samples of SNIA (Ref. 65). Though possible non-monotonicities in the Hubble diagram were already investigated in Ref. 63, no significant wiggle was detected. These might lead to greater effects than the small scalar perturbations in the Robertson-Walker metric of Ref. 64.

But the important result is that such scenario makes clear why we must retain the t_{clock}^α mean regime resulting in this succession of alternate t_{clock}^2 and $t_{clock}^{2/3}$ for cosmology. The power law t^α with α between 1.52 ± 0.11 (SCP 2008) well fits the Hubble diagram up to redshift 1.8. Any α can be obtained depending on the relative time elapsed in the two regimes. The power law obtained from the SN suggests that our side has been mainly in the constantly accelerated regime appart from small excursions in the $t_{clock}^{2/3}$ regime.

However we should rather wait for the nearby SN factory results to get confidence in the Hubble diagram method, since the physics of discontinuities might be involved in the processes leading to the Supernovae explosions and this in turn directly depends on the scale factor behaviour. Such an unexpected evolution mechanism might complicate the extraction of the cosmological information from the SN Hubble diagram.

18.2. Nucleosynthesis

BBN theory predicts the universal abundances of the light elements D, He^3 , He^4 and Li^7 which are essentially determined by $t \approx 180s$ (Ref. 55). The best observable to probe these predictions is the abundance of He^4 relative to H. The mass fraction $Y_p \approx 1/4$ is tested at a ≈ 5 percent level of accuracy and is very sensitive to the expansion rate at the time of primordial nucleosynthesis. A two orders of magnitude smaller expansion rate would result in a negligible abundance of Helium since there would be much more time for the disintegration of neutrons between the time when the neutron-proton inter-conversion rate were frozen by the expansion and the later time when nuclear reactions could proceed to incorporate protons and the remaining neutrons into light nuclei. In DG, because the expansion rate extrapolated at $10^9 K$ is orders of magnitude lower than in the standard model because of the accelerated or constant ($\alpha \geq 1$) expansion rate, it thus seems hopeless to get a non negligible Y_p . However, as we could check with in simulation, a kind of miracle happens: with an expansion rate many orders of magnitudes below the SM one, Y_p rises and reaches $Y_p \approx 1/4$ again. Authors have investigated the possibility of having the good proportion of Helium in universes expanding much faster than in the SM and found that for an exponent in the power law greater than 1, it is indeed possible to obtain the correct Y_p with a baryon to photon ratio significantly different than in the Standard Model (Ref. 60 and references therein).

For instance, Figure 2 in Ref. 61 shows that $\alpha = 1.52$ gives the correct Yp for a density of baryons at the level of the percent of the critical density! Since this is in perfect agreement with the density of visible (actually detected) baryons we don't even need to explain an important deficit (≈ 1 percent is about one fourth the baryon density predicted in the standard model) in the form of cold molecular hydrogen dark clouds as is necessary in the standard model.

Let us notice that $\alpha = 1$ gives the correct Yp for a density of baryons an order of magnitude greater than that of the Standard Model. However, whatever α , the baryon fraction just after BBN could be considered an upper bound for the present one since an unknown fraction of them might have annihilated with antibaryons from the conjugate side between BBN (most probably just after BBN) and now.

If DG is successful in this confrontation, we must conclude that the SM ability to predict the good abundances was a pure coincidence. Let us list the reasons why this coincidence is not as improbable as it might appear at first sight. Indeed the only two other observables on which BBN rests up to now are the D/H and Li^7/H fractions. The systematical effects are at the level of the measured value for Li^7/H so it is not clear at all whether we are here sensitive to a primordial abundance. Unfortunately, the same conclusion applies to D/H since its systematics (dispersion between measures) are not understood, in particular the D/H large inhomogeneities, a very serious anomaly in the absence of any known astrophysical mechanism to explain either the creation or the depletion of D (Ref. 54). The physics of discontinuities is likely at the origin of these anomalies since nuclear transmutations are commonly observed in discharge experiments in relation with the presence of micro lighting balls that represent the most perfect signatures of a spherical discontinuity. See for instance Ref. 86, Ref. 87 and Ref. 85.

18.3. *Structure Formation and the Early Universe*

We here assume the case of an homogeneous and $t^{4/3}$ accelerated universe. The qualitative conclusions would remain the same for any α between 1 and 2. Taking the dominating mass density contribution to be the baryonic matter density ($\rho_0 = 0.03\rho_c$ given by the nucleosynthesis constraint from the previous section we get at the time t_R of hydrogen recombination the density through:

$$\begin{aligned}\rho(t_R) &= \rho(t_0) \left(\frac{a(t_0)}{a(t_R)} \right)^3 \\ &= \rho(t_0) (1 + z_R)^3 = \rho(t_0) (1500)^3\end{aligned}$$

Also, we can link the Hubble parameter at t_R and nowadays value at t_0 through:

$$\frac{H(t_0)}{H(t_R)} = \left(\frac{a(t_R)}{a(t_0)} \right)^{3/4}$$

thus:

$$H(t_R) \approx H(t_0) (1500)^{3/4}$$

Neglecting the effect of the universe expansion in the evolution equation of density fluctuations $\delta(t)$ on our side and making use of $p \ll \rho$ after recombination leads us to the following differential equation:

$$\ddot{\delta} - 4\pi G\rho\delta = 0$$

and the exponentially growing fluctuations:

$$\delta_+ = e^{\sqrt{4\pi G\rho}t}$$

Then we can check that:

$$\frac{H(t_R)}{\sqrt{4\pi G\rho(t_R)}} = \sqrt{\frac{2}{3}} \frac{1}{1500^{3/4}} \sqrt{\frac{\rho_c}{\rho_0}} \approx 2.10^{-2}$$

insuring that the universe expanding rate was indeed negligible compared to the density fluctuations growing rate thereby justifying our previous approximation. This accelerating universe from the CMB last scattering up to now is older than the standard model universe: $4/3H_0 \approx 18$ billion years.

But our universe is also much older at the time of Hydrogen recombination than it was usually believed to be in the Standard Model framework:

$$t_R = \frac{4}{3H_R} = t_0 \frac{H(t_0)}{H(t_R)} = 18.10^9 \left(\frac{1}{1500} \right)^{3/4} \approx 75.10^6 \text{ years.}$$

provided we can neglect the earlier exponential regime duration. At t_R , the universe expanding rate was already (it decreases in time) so small that it did not affect at all the growing of primordial fluctuations so that we could reach soon the non-linear regime starting from the 10^{-5} density fluctuations of the CMB. We notice also that the smallest mass fluctuation able to grow after recombination for the present universe density $\rho_0 = 3.10^{-31} \text{ g/cm}^3$ is the Jean mass $\approx 10^8 M_\odot$ (see in Ref. 9 the plotted Jean Mass as a function of temperature for different values of the density) while the typical dwarf galaxy baryonic visible mass is $\approx 10^9 M_\odot$.

The larger dimensions of voids interpreted as structures in the conjugate form indicate an initial asymmetry between the two conjugate universes content as first suggested by JP Petit (Ref. 1) whose simulations have confirmed the appearance of large voids with galaxies concentrated at the periphery of these repelling conjugate structures. The matter/anti-matter symmetry was satisfied at the origin of time when the two sides of the universe merged. After separation, the annihilation of antimatter and matter on both sides must have produced as much matter remaining on our side than antimatter on the conjugate side and an a priori different density of radiation but statistically negligible relative difference $\propto \frac{1}{\sqrt{N}}$ considering the large number N of annihilating particles involved. Therefore the density of baryons and photons are the same on both side. The origin of the asymmetrical evolution is obvious : if our side mainly evolved according the t_{clock}^2 regime, the conjugate side must have mainly evolved according the $t_{clock}^{2/3}$ regime. This is probably the main reason for the nowadays very large asymmetry between the typical sizes and shapes

of structures on the conjugate sides. In this regime our clocks are also accelerating with respect to the conjugate side clocks so that the conjugate side may be somewhat late and in particular, its matter-radiation decoupling and formation of structures probably started later.

Therefore, we found that our model is not only successful in explaining the growing of the smallest initial CMB fluctuations in the linear regime without any need for dark matter nor dark energy but also could lead to an efficient formation of galaxies and universe voids interpreted as over-densities in the conjugate form. In this derivation the very small expanding rate in the early universe played a crucial role.

18.4. *Primordial fluctuations and the CMB*

Most of the CMB signal is clearly the signature of a hot and dense primordial state of the whole observable universe. Its most remarkable properties are its black body spectrum and its homogeneity far beyond the horizon predicted by the standard model at decoupling. For this reason the inflationary scenario has been proposed to save the model. It is therefore important to notice that the DG universe evolution started from an almost stationary regime and has been accelerated ever since. Consequently it has maintained a much smaller expansion rate than in the SM in its earlier ages and was already much older than the SM universe at the time of decoupling so there were enough time to let the fluid become homogeneous over a much greater scale than that defined by the Standard Model Horizon scale at decoupling. In the absence of any horizon mechanism to explain the scale associated to the peaks of the CMB spectrum, our context is very different from the Standard Model one (Ref. 24). We would be in trouble to obtain the right position of acoustic peaks relying on the standard model mechanism (the attempt for a freely coasting cosmology is in Ref. 67). However recent studies show that the cosmological constant assumption is essential in the framework of the Λ CDM model to reach the percent level of accuracy on the curvature. Giving up this hypothesis, a more reasonable 15 percent error on the curvature parameter (Ref 82) is obtained. This along with various claimed anomalies of the CMB especially at the largest scales but also the too many parameters entering in the CMB spectrum fit allow us to seriously doubt that the standard model and inflation actually predicted the position of the first peak or the correct shape of the CMB spectrum.

In our framework, the fluctuations we now see in the CMB spectrum are most probably not related to properties of the CMB at decoupling since we have no obvious mechanism to predict the observed scales of the peaks. Rather the explanation has to be searched for in the inhomogeneous universe in between last scattering and now. By the way, let us mention that DG does not even need detectable primordial inhomogeneities to insure the subsequent formation of structures thanks to the almost exponential growing of any inhomogeneity after decoupling in a very slowly expanding universe.

Looking for a mechanism that could affect our side CMB photons wavelengths at a relative level of 10^{-5} at the one degree angular scale during their path from the surface of last scattering to our detectors, we cannot rely on the usual ISW effects because they are absent in DG. As for the SZ effects, their contribution is only significant on much smaller scales than the degree. DG discontinuities and Gravitational waves also cannot inhomogeneously affect our side CMB photons (in a $B=-A$ field). The solution comes from the fact that the CMB radiation is also present on the conjugate side and that before the conjugate sides separated the photons fluid were common to both sides. Therefore the CMB photons from both sides should appear redshifted by cosmological expansion by exactly the same amount at a given time from the point of view of an observer on our side though from the point of view of an observer on the conjugate side both must appear much hotter. The important point is that at any time from our side point of view the CMB is present on both sides of the universe at the same temperature apart from fluctuations which origin we are going to explain and suggest now in relation to local $B=-1/A$ field inhomogeneities.

Our suggestion is that inhomogeneities of the CMB are produced when the CMB radiation from the conjugate side appear on our side via discontinuities that we expect to encounter more frequently in the vicinity of various foregrounds : indeed structures at any scale generate potential wells and the deeper the potential wells the lower the drifting speed of the discontinuities in their vicinity hence the longer the mean time such discontinuities will remain in these regions as we explained earlier.

When CMB photons from the conjugate side cross the gap and appear on our side for instance at the bottom of a deep potential well, their frequency is shifted relative to the frequency they had on the corresponding hill of the conjugate side. This shift does not seem obvious because photon frequencies are never affected in a stationary gravitational field whether it is discontinuous or not. Indeed for instance the usual gravitational redshift is just a comparison of two clock rates at different gravitational potential levels, comparison made possible precisely because photons retain the frequency at which they were emitted during their path from one clock to the other. But when they reach a discontinuity and cross the gap, photons are not understood to be in a stationary gravitational potential there but in a field which time $\tilde{t}$ reverses into $t=-\tilde{t}$. This reversal might be thought of as a continuous complex rotation (thus involving an extra dimension) from $\tilde{t}$ to $t = e^{i\theta}\tilde{t}$ where $\theta = \pi$. During this time reversal the field $\tilde{g}$ is transformed into $g=1/\tilde{g}$ thus the photon frequency is shifted by twice the local gravitational potential. Therefore, contrary to the case of photons staying on the same side which are never affected by local potentials they travel through (in part because there are no ISW effects in DG) photons crossing the gap from the conjugate to our side are shifted. The peculiar galaxy velocities (1000 km/s) tell us about the typical level of the potential wells at work in the universe $Gm/rc^2 \approx v^2/c^2 \approx 10^{-5}$: exactly what we need to account for the amplitude of the CMB temperature fluctuations. It is also encouraging that

a recent analysis (Ref. 100) has shown that cosmic strings may account for roughly 10 percent of the CMB map of fluctuations. These may rather be the first signature of the conjugate side CMB shifted photons emerging on our side through cosmic string like discontinuities.

The transfer of a photon from the conjugate to our side is not understood to be an absorption followed by a reemission by a discontinuity. When the two sides background fields cross each other the wave function of a photon can split at the discontinuity and a QM computation should allow to compute the probability for the photon to remain (quantum collapse of the wave function) on the same side or alternatively be transferred to the conjugate side. In neither case do we have the extra expected double Doppler effects produced by absorption and reemission of a particle. This is fortunate because the discontinuities are in motion in the universe and their speeds relative to the observer would produce much larger temperature fluctuations just due to the Doppler effects than what is needed to account for the CMB fluctuations amplitudes.

If the CMB foregrounds are various structures with the needed discontinuities in their vicinity responsible for shifting the frequencies of conjugate side photons and in this way producing all apparent CMB fluctuations, let us examine if it is possible to obtain the correct typical 1 degree angular scales from known structures. While the large and small magellanic clouds in the vicinity of our galaxy and Andromeda correspond to 10, 3 and 2 degrees apparent angles respectively, the number of galactic objects really explodes at angles smaller than 0.1. Neighbour clusters of galaxies with their typical apparant 1 degree or subdegree scales which is also the scale of the typical distances between them are thus interesting candidates to account for CMB power spectrum at this scale. The repartition in the sky of these more distant objects is also naturally much more homogeneous but how could we explain peaks in the spectrum. Very interestingly there are evidences, even in the more recent works, of cluster concentrations at particular redshifts, roughly multiples of 0.03: $z=0.03$, 0.06, 0.08, 1.1 corresponding to apparent sub-multiples of an angular scale about 1 degree corresponding to 2Mpcs diameter clusters at $z=0.03$. Roughly the same scale appears in the typical distances between those clusters. However we would expect a clear correlation between the distribution of these clusters and CMB maps if these were the main contributors. Instead recent estimation of the ISW effects (Ref. 101) have shown that the correlations are much lower (even lower than the SM predictions for the ISW effect) than needed to establish a link with the known amplitude of the CMB main peak.

But there is may be an even more interesting candidate: the scale of the superclusters and supervoids of the universe (Ref.102) because at this scale a quasi-periodic feature in the repartition of these superstructures would hopefully result in a succession of harmonic peaks in the CMB power spectrum. These structures are seen under a four degrees apparant angle at $z=0.35$ (this is our alternative interpretation of the so-called BAO peak) and we compute that in DG the corresponding

angular scale was that of the first CMB peak at $z \approx 8.4$ (the age of reionization) rather than $z \approx 1100$ in the Standard Model. It might be that the discontinuities left their imprint on the CMB at this epoch.

At last, a recent study has shown what might be considered as a kind of honeycomb (Ref.103) structure in our own galaxy, interstellar clouds delimitating empty bubbles (rarefied gas) with a typical diameter of 150 pcs, like our own local bubble centered on the solar system a new map has revealed. Such periodical pattern would arise in the CMB spectrum at the one degree scale if its contribution to the fluctuations is the greater at a distance of nearly 9000pcs (comparable to the distance between the sun and the galactic center) from us in all directions. The "cold fingers" joining the galactic pole in WMAP maps also suggest a local contribution.

Now what about the contributions on the largest scales of the lower multipoles? The most important energy in the quadrupole that was found in a suppressed foreground (Ref. 78) of WMAP, a restricted area corresponding to the direction toward the galactic center is almost the perfect signature for our scenario. This is sufficiently powerful to indicate an extra source of CMB foreground in the direction of the largest concentration of stars (i.e. many potential wells thus discontinuities) in our vicinity: that of the galactic center. The lack of energy at the highest scales would simply be the consequence of the concentration of our galaxy stars in a plane, resulting in an easily removable foreground, just by applying a suitable mask.

A local, anisotropic with respect to the earth observer, discontinuity in the solar system would ideally account for the correlations with the equinoxes directions and the ecliptic plane seen in the CMB low multipoles by WMAP (see Ref. 23) but a systematical effect or software error is not excluded. Indeed the Wmap data have recently been re-analyzed and the authors claim that they succeeded to have their quadrupole almost completely removed by better correcting Doppler effects depending on the WMAP spacecraft attitude parameters (see Ref. 94, Ref. 95 and references therein). May be this is the better confirmation we could hope for that there is no cosmological signal in the lowest multipoles corresponding to scales concerned by the foregrounds which are the most easily removable simply by using a mask. This reinforces our conviction that the cosmological fluctuations signal would disappear at all scales if we could remove foreground effects at all scales. Even the Wmap Ref. ? collaboration admits that there also exists a kind of quadrupolar modulation of all the multipoles (detected at a 9 sigma significance level) that most probably is related to a systematical effect since it is strongly correlated with the direction defined by the ecliptic plane.

The resemblance of the WMAP power spectrum with vanishing lower multipoles to the spectrum of an AMI bipolar (or order 2 bipolar) random signal would also be striking. In potential wells (resp hills) where the discontinuities are more numerous we would get the most important fluctuations (bit 1 of the AMI signal) from alternating zones of redshifted (resp blueshifted) CMB photons with AMI polarity=+1 (resp AMI polarity=-1) and unberturbed zones in between (bits 0 of the

AMI signal) far away from the structures on both sides.

Many other possible consequences of discontinuities, in particular in our solar system, were explored in Ref.79.

A recent publication (see Ref. 89) using the WMAP 3 years data gives a very satisfactory description of the CMB spectrum on the largest scales by postulating the topology of a flat torus for our universe. An $L=3.86$ Gigaparsecs torus gives our universe a finite volume of roughly $5.10^3 Gpc^3$ but can explain the spectrum at scales larger than the Horizon scales without the need for inflation simply because by generating the optical illusion of the infinite repetition of this pattern, it produces higher wavelength contributions. The scenario is very interesting for us because it proves that a repetitive pattern on the CMB can lead to a succession of peaks which is the idea we rely on to get the main CMB oscillations starting from the one degree peak. These regularities might be the signature of a universe structured according a periodical network and one should be aware that such regular pattern may be apparently lost beyond some distance because of the particular angle at which the network is rotated with respect to the observer line of sight. A periodical structure would be an important confirmation of another prediction of DG: the structure of vacuum according a periodical network which would have left very small fluctuations (invisible) in the primordial plasma, at the origin of the nowadays, after expansion, 100Mpc typical cells. Of course any network also manifests preferred directions and this is why some of the most intriguing CMB properties are very promising for us (See the recent Ref. 91 that seems to confirm the existence of preferred directions).

18.5. *Galaxy Rotation Curves*

The subsequent structures nonlinear evolution is also facilitated given that an older universe $(4/3)H_0 \approx 18$ billion years from the CMB last scattering up to now, to be compared with the oldest ($z=5$) galaxies ages approaching 13.5 billion years, in quite a good agreement with the oldest stars ages, provides more time for galaxy formation. Notice by the way that our model is not in trouble with the oldest stars and recently observed old galaxies as is the Standard Model with the new very accurate estimation of the standard universe age by WMAP. Though this should be checked in details with numerical simulations, it appears already almost certain that the interaction between conjugate density fluctuations is all we need to solve missing mass issues at any scale. The problem was already addressed in some details in the work of JP Petit, particularly in Ref. 1 where he could show that conjugate matter is a powerful alternative idea to dark matter or dark energy. In DG, the repelling effect of a galaxy in our metric must generate a large void in the conjugate universe. This in turn is completely equivalent to a huge halo of perfectly homogeneous matter source density on our side. This halo is of course completely dark and only interacts gravitationally with our galaxy. Because similar halos of weakly interactive dark matter can help to obtain satisfactory galaxy flat rotation curves, our model is very promising. The main difference with the Halo of

weakly interactive dark matter is the perfect homogeneity of a void which prevents any small scale gravitational collapse to enter into the game. Very remarkably, it has been shown in Ref. 22, that a model which suppresses gravitational interactions (GraS) between dark matter and baryons on small, subgalactic scales (1 kpc) allows to get the correct inner rotation curves of small galaxies and avoid the formation of too many substructures in galactic halos. The issue is that the CDM Standard Model, though it is very successful on the large scales predicts very compact density cusps in the Dark matter halo and significantly more object of dwarf galaxy mass in a typical galaxy halo than what is seen. Both predictions are related to the gravitational effect of DM aggregates on small scales but conflict with observations. In our framework, the gravitational artificial suppression is not needed since the smooth central potential profile is a natural consequence of the void nature of the conjugate fluctuation (nothing concentrated at the galaxy center!). Moreover, as in the GraS model, baryons here do not fill dark matter substructures simply because the latter do not exist in the void of the conjugate metric. The recently announced discovery of Dark Matter by the NASA in Ref. 43 is a bad new for MOND and TeVeS since these theories rely on a modified gravity to account for the rotation of galaxies. This is not the case in DG which in the weak field approximation is as Newtonian as GR. DG also has Dark Matter but it gravitationally repels our side matter and the Chandra X ray data will have to be examined in this perspective as any other data sources. A shell of Dark repelling matter may be very efficient to help obtaining a bar spiral galaxy stable structure as shown in the 2d simulations by JP.Petit and F.Landsheat Such shells were predicted in the work of Ref. 1. One of the most impressive indicator that there actually is a fundamental scale of mass surface density at which the gravitational accelerations suffer a drastic change in regime is the ability of MOND theories (81) to accurately fit the measured galaxy rotation curves by switching to a modified gravity when accelerations fall below a fundamental threshold. In our framework it is very tempting to identify this density as being the vanishing one resulting from the perfect compensating positive and negative densities from the conjugate sides of the gravitational field. A recent simulation result trying to mimic the good performances of MOND fits through an ad hoc distribution of DM can, according its authors, both reproduce our galactic rotation curve and observed diffuse galactic gamma rays excess by EGRET (see Ref. 52 and references therein). Along with the usual distribution of Dark matter, this study assumes the existence of two rings of Dark matter in the galactic plane at 4kpc and 14kpc where Susy Wimps annihilation produces the observed gamma ray excess. But Ref. 74 shows that this model is excluded by a wide margin from the measured flux of antiprotons. DG would quite differently try to fit the data. In our framework, we would need shells of the conjugate form anti-matter to produce both the gamma rays and some modulations in the galactic rotation curves provided, as in the vicinity of our galactic center where positrons excess were observed, antimatter from the conjugate form can cross the gap toward our form and annihilate with normal matter near these rings. Obtaining rings with antimatter would also be

much more natural since it is not collisionless as is Dark matter. The stability of antimatter shells would be insured because these would be surrounded by normal matter which is repelling from their point of view but since these shells repel in turn our matter they should be located at somewhat different places than the DM rings of Ref. 52 to get the correct deformation of the galactic rotation curves.

18.6. Large Scale Dynamics

At last, we know that the standard CDM model is able to correctly predict clusters abundances and galaxy peculiar velocities. Since our framework allows us to take into account the effective large mass induced by the dark side voids, our galaxies are much heavier and extend very far away from the visible part which in some case might be too faint to be detectable. Because again the equivalence between the CDM dynamics and our dark gravity with voids dynamics seems natural, successes of the CDM model should convert in a straightforward way into our model success on the large scale provided there is no bad surprise from more detailed simulation studies.

19. A cosmological scenario to satisfy the Baryonic oscillation constraint ?

The baryonic peak was discovered (Ref. 90) recently in the spectrum of galaxy clusters at a redshift about 0.35. The general opinion is that it corresponds exactly to the CMB main peak at 1 degree in the sense that we are dealing with exactly the same scale which were only submitted to the effects of cosmological expansion between $z=1100$ and $z=0.35$. The ratio between the angular distances to the BAO ($z=0.35$) peak and to the first CMB ($z=1100$) peak was measured to be $R = 0.0979 \pm 0.0036$ and this is believed to be a strong constraint. This constraint is obviously not satisfied by our accelerating solution which would prefer a much smaller R . However, it is not obvious that the two observables are directly linked to each other. Indeed the 143 Mpc "baryonic oscillation" peak scale is very close to the typical scale of the supervoids in the universe (133 Mpc for the diameter of the included sphere) and the strange scale of more intense galaxy clustering at distances multiples of 180 Mpc ($z=0.03$) which both have nothing to do with the CMB first peak. Our privileged interpretation is that the 3 length scales are related to each other since they correspond to distances in a network structure. It might be that this structure left its imprint on the CMB at $z=8.4$ which would produce the first CMB peak at the observed angular scale.

20. Precession of Equinoxes and GP-B Gyroscopes

In the context of a theory involving discontinuities, it is worth reconsidering the old question that opposed Newton, Mach and Einstein of what should determine the inertial frames. In the spatial volume enclosed within a discontinuity, let us call this a bubble, we suspect that there are associated globally inertial frames,

where globally here means valid everywhere inside such bubble : a globally inertial frame is a frame where inertial pseudo-forces (but in general not gravitational forces) vanish everywhere in the bubble. Usually such frames are understood to be not rotating but not rotating with respect to what? Newton would have answered with respect to absolute space, Mach would have answered with respect to the fixed stars, Einstein would have answered with respect to the total gravitational field generated by local sources plus global background. The presence of a discontinuity that isolates some part of the universe from all the rest of it, leads us to another likely answer. In DG, a globally inertial frame of a given bubble is one which is not rotating with respect to this bubble but in general possibly rotating with respect to the fixed stars. If such frames exist different from the universe restframe, the expected phenomenological consequences should be simple: any gyroscope axis (earth, earth-moon system, earth-sun system, Gravity-probeB gyroscopes) within the same bubble should follow the rotation of the globally inertial frame associated to this bubble.

It is well understood that the differential gravity forces from the sun and moon on the earth oblate spheroid are responsible for the precession of the earth spin axis with a well established 25580 years periodicity, which translates into the annual precession of equinoxes of $360.3600/25580 = 50.3''$ per year. If the precession frequency of the earth natural gyroscope was somehow determined by the rotation of a local bubble with respect to the universe restframe rather than by these tidal earth-moon effects, then the relative position of the earth moon and sun would also have been stabilized just in such a way as to have the frequency of the tidal induced precession rate matching $50.3''$ per year in a resonant way. The motivation for seriously considering this scenario in DG is obvious, since for the solar system we have a discontinuity scanning the whole solar system precisely with the same period (at the 15 percent level of accuracy) determined earlier from the Hubble constant H_0 and solar surface potential. Hence, we are just postulating that the discontinuity path period is also that of a rotating frame associated to a bubble encompassing the whole solar system. The consequences are obvious for the four Gravity-ProbeB gyroscopes which should exactly mimic the deviation of the earth spin axis: thus a $50.3''/\text{year}$ equinox like precession which is hopefully the effect (it has the same magnitude) the collaboration described as a classical electromagnetic effect. By the way, the same globally inertial effect on WMAP and COBE satellites might be responsible for a drift in the CMB Dipole apex which would also take the same value of $\approx 50.3''$ per year between COBE and WMAP1. If Ref. 77 is correct the effect, though not yet statistically significant between COBE and WMAP1 might already have been observed between WMAP1 and WMAP7.

21. The Discrete Mode of Vacuum and the Origin of Mass

21.1. *The elementary source and field*

As we already noticed earlier, we have on the source part of our DG B=-1/A field equation ρA rather than ρ and the resulting effect is not exactly balanced by gravific pressure effects as it is in GR. $A = e^a$ in the source term clearly stands for the self gravitational potential energy of the source and its gravific effect. This can only be completely understood by solving the equation satisfied by $a(r)$ inside matter, which happens to be the Liouville equation (still in units such that $c=1$):

$$\Delta a = -8\pi \bar{G} \rho e^a$$

where we anticipate that $\bar{G}$ is not necessarily G but must be unambiguously related to the familiar G constant. Let us assume that nothing but a spherical source with constant density ρ , radius R hence energy $M = \rho \frac{4}{3} \pi R^3$ is the elementary source for $a(r)$.

The solution which appears to be unique inside the source is $a_{in}(r) = -Ln(4\pi \bar{G} \rho r^2)$ or $a_{in}(r) = -Ln(3\bar{G}M(r)/r)$ where $M(r)$ is the total energy within radius r . Of course the solution outside the source of $\Delta a = 0$ must be $a_{out}(r) = 2GM/r$ if we want to recover the familiar Newtonian potential in the weak field approximation as well as the asymptotic Minkowskian behaviour of the field elements B and A . But at r equal R , $a_{in}(r)$ and its derivative must match $a_{out}(r)$ and derivative. $a'_{in}(R) = a'_{out}(R)$ can only be satisfied provided $GM/R = 1$ i.e. the elementary source radius must also be its Schwarzschild radius. Then $a_{in}(R) = a_{out}(R)$ can only be satisfied provided $2 = -Ln(3\bar{G}/G)$ or $\bar{G} = G/(3e^2)$.

The condition $GM/R = 1$ is certainly one that cannot be satisfied by the vast majority of known extended sources as we understand them usually and yet our solution seems to be the only valid one we can propose in DG in case of spherical symmetry and constant density inside our source. However remember that our B=-1/A field is only the field generated by what we have called an elementary source and it is tempting to identify our elementary source with an elementary particle which indeed might be conceived as such a micro object with Schwarzschild radius just equal to its radius (this might even be applied to a photon if it has even an extremely small rest energy). With this understanding of elementary sources, as we already explained earlier we recover all predictions of GR including gravitomagnetic and pressure effects after exporting all elementary solutions and combining them in any common coordinate system, except of course the departure from GR at the PostPostNewtonian order of our Schwarzschild solution.

However there is an even much more attractive possibility which we are going to present and explore now, the possibility that the mechanism that relates $T^{\mu\nu}$ to $S^{\mu\nu}$ might involve a discrete mode of vacuum, a physical ether made of a genuine network of masses in gravitational interaction (worth reading are Ref. 71 and Ref. 58 though i'm not sure the author's idea is the same as that explained here). We postulate that what we usually call vacuum is actually a physical entity, a genuine physical

ether which mass is not continuously distributed over space but is concentrated in a discrete network of the kind of masses our previous investigation of the $B=-1/A$ equation led us to: each mass m_0 has a Radius R_0 equal to its Schwarzschild radius Gm_0 . Of course the gravitational stability of such a network is only granted provided positive and negative masses alternate in such a way that any mass has only repelling closest neighbours in its vicinity. Even the observational effects expected from this background of huge alternating masses may not have been evidenced so far because of the globally compensating positive and negative gravitational effects particularly if these are separated by small distances d in the network and if we happen to be ourselves in motion at high speed with respect to the restframe of this network. Now because the needed objects satisfying $R_0 = Gm_0$ are already there everywhere in the vacuum and because we also have in DG discontinuities which if located at radius R could explain why some rest energy can indeed be trapped within the sphere of radius R we can come back on and strongly support our earlier intuition that gravific energy i.e. mass is only generated by such objects and gives rise to $S^{\mu\nu}$ which only non vanishing component is S^{00} : the total energy trapped by the spherical discontinuity at $r=R$. Whenever a wave packet of any field propagates in the network, some of its energy E (a quantum as we shall argue later) can be trapped (absorbed) then reemitted by such objects which radius R vary accordingly to maintain the condition $R = G(m_0 + E)$. Only in this way can the energy of the field source gravity while the energy it carries in between the Masses m_0 of the network does not source gravity : of course it cant because the conditions required by the $B=-1/A$ field equation are not satisfied for this to be possible.

Eventually, we should be aware that by solving the elementary field equation we already took into account self gravitational potential energy effect of the $B=-1/A$ field itself and simply recovered our beloved solution $g_{00} = e^{-2\frac{Gm_0}{rc^2}}$ outside the elementary source where m_0 is only $\rho\frac{4}{3}\pi R_0^3$. However, remember that ρ itself does include also the the gravitational potential energy of our source in the total external field of all other elementary sources. If in a compact star, there is a single unique privileged frame, for instance the restframe of the star where we are allowed to compute a myriad of elementary $B=-1/A$ fields then combine them, this will result in a total field that will really take into account the gravific effect of the self gravitational energy of the star in its own field but not the pressure effects and as we stressed earlier this can have very important observational consequences such as the generation of a corrected gravific mass $m_0 A_{ext}$ which can be huge if A_{ext} is strong.

With our expression for $g_{00} = e^{-2\frac{GA_{ext}m_0}{rc^2}}$ in the freefall equation, one can derive the following exact general expression for the gravitational potential $\frac{c^2}{4}(e^{-4\frac{Gm_0 A_{ext}}{dc^2}} - 1)$ created by m_0 which in the weak field approximation gives back the Newtonian potential $-Gm_0 A_{ext}/r$.

21.2. *The network and its vanishing energy conditions*

The $B=-1/A$ field is only generated by density perturbations relative to a mean homogeneous density of the universe. Thus if the vacuum mass was homogeneous, there would be no associated gravitational potential energy of a single newly created mass at rest in the $B=-1/A$ Minkowskian field generated by the universe vacuum. The energy needed to create this mass m_0 would simply be the sum of its rest energy and the total integrated energy over space of the $B=-1/A$ gravitostatic field it generates which t_{00} we computed earlier is $\frac{-2a'^2}{16\pi G}$.

Using $m(r > R_0) = m_0$ and $m(r < R_0) = m_0(r/R_0)^3$. The total integrated t_{00} then yields $-12e^2m_0$. As a consequence, the creation of such a single mass at rest on one side of the Janus universe cannot be at free cost because $m_0 - 12e^2m_0$ does not vanish.

The only possibility to explain the generation of energy and radiation starting from an empty universe is a theory allowing rest mass to be created at free cost. Then the subsequent decay of such primordial rest mass into less massive particles could transfer part of this rest mass energy into kinetic energies and therefore all forms of the energy could be generated in the universe starting from nothing (at free cost).

But fortunately a vanishing energy condition can be satisfied by each mass m_0 in our vacuum just because its mass is not continuously distributed over space but is concentrated in a discrete network of such m_0 masses. Indeed the vanishing total energy condition for each mass must also include its gravitational potential energy in the total field generated by all the other masses.

For instance in case we would just want to create a pair of opposite masses m_0 at a distance d from each other, one on each side of an otherwise empty universe, the vanishing energy condition is satisfied provided the distance d between them is constrained by (avoiding double counting of the potential energies and neglecting A_{ext} effects):

$$m_0 \frac{1}{4} (e^{4\frac{Gm_0}{d}} - 1) \frac{1}{2} + m_0(1 - 12e^2) = 0$$

Playing the same game but taking into account all other masses in an infinite network, the vanishing energy condition would involve

$$\left[\sum_i \frac{\pm 1}{d_i} \right]$$

in place of $1/d$. With alternating signs from a mass point to its neighbours in a cubic network, the serie converges to $1.698/d$ in a cube of side L when L tends to infinity so that eventually we get a simple relation linking d to m_0 thus R_0 .

$$\frac{R_0}{d} = 0.96$$

thus d is a little bit greater than R_0 which insures that two neighbour masses of the network on the same side of the universe are separated by $2d$ hence do not overlap.

It's less a concern that actual neighbour masses are separated by d less than $2R_0$ because these are not on the same side of the universe. Notice that since at such distances we are in the strong field regime, A_{ext} effects should have been taken into account, however this treatment is a priori difficult so we let it as an open issue for the time being. Then the important result if confirmed would be that eventually the network of mass points can be created at free cost with masses m_0 in spheres of radius R and corresponding step distances d between them satisfying a simple relation.

The network topology must be well constrained. Indeed, the simplest polyhedron to pave a flat space is a prism with two equilateral triangle faces and three square faces. However it does not allow the alternate mass signs from one point to any neighbour which is necessary to insure the gravitational stability of the network. A Cubic network or a network paved with hexagonal based polyhedrons or even Penrose networks are particularly interesting among other cristallographic possibilities.

We saw that the DG B=-A sector has massless longitudinal wave solutions. Extending this framework to include extra dimensions, the massless waves can acquire mass from the point of view of our 3+1 space-time provided the extra-dimensional path is quantized. This is just the result of the usual assumption of a compact extra-dimension which length scale l determines the lower mass of the Kaluza-Klein tower of states. We can alternatively consider a non compact extra dimension but the fraction of our wave momentum in the extra dimension would still be quantized if the wave is periodically created and annihilated at the network points in such a way that its trajectory is zigzagging in the extra-dimension, its momentum component in this extra dimension being transverse to the mean observed momentum. This "missing" transverse momentum might generate a mass and a complete spectrum would follow from the geometrical properties of the network (characteristic lengths and angles) in the 3+extra spatial dimensions.

The idea is attractive but unavoidably amounts to admit fundamental massive fields in our 3+1 dimensions while we earlier explained why our framework instead favours fundamental massless fields with the left and right fields being respectively restricted to our and the conjugate side of our universe.

Without appealing to extra-dimensions, the interaction of fundamental massless fields with our ether, the vacuum network of mass points, might be another possibility to generate an effective mass (why not an entire spectrum?) for these fields just in the same way as photons acquire an effective mass propagating in a transparent medium. But then also Lorentz violating effects associated to the privileged frame of this ether are expected and it would hardly be explained why such Lorentz violating effects remained unobserved so far. This however remains an option to generate very tiny thus still unobserved effective mass terms but hardly the masses of the standard model fundamental fermions.

Other options to get mass? We earlier already expressed our frustration as regards spontaneous symmetry breaking or other theoretical ideas that just help to understand the generation of mass in our universe but up to now are not able to

predict the actual mass spectrum starting from a reduced number of fundamental constants.

Eventually remains the possibility that fundamental fermions be massive composite objects made up of bound states of fundamental massless fields. This is nearly the case for the proton which mass is essentially determined by the gluons interactions rather than the masses of its quark components.

By the way, let us mention that this is an intriguing result and interesting from our framework point of view. Indeed if the treatment of a bound system as a source for gravity was the same at this small scale of the nucleon as it is at the scale of a star, one would expect the gravific mass of the nucleon to be determined by the mass of the quarks, the other contributions, i.e. potential energies and internal pressure terms (pressure of the partons inside the nucleon) cancelling each other just as pressure and gravitational energy potential terms do in a static configuration of a star (see again Ref 28). In other words the equation of state for cold baryonic matter could not be $p \approx 0$ as usually assumed! Such an observation is clearly favouring a framework where pressure terms do not contribute to the gravific mass: just what we have in DG!

22. Quantizing Gravity

We have introduced a network of vacuum masses. The next step is to realize that even if gravity is only sourced by the energy density ρ inside a sphere of volume R_0 , the actual field which energy density is ρ there does not need to be confined within the sphere. It can actually extend much beyond and even overlap with other fields. The unique characteristic responsible for this particular field to be a possible contributor to the gravific ρ inside the sphere is that it is a spherical field which center of isotropy is the center of the sphere. Since the total energy of the field is not trapped inside the sphere, actually it's just a fraction of this energy that is sourcing gravity, the exact value of this fraction being determined by the value of the gravitational constant G .

So we can now consider that this field has a total energy which is not trapped inside the sphere of radius R_0 but rather by a spherical discontinuity at a radius R much larger than R_0 . Therefore the field can only be trapped in the form of a superposition of standing waves : the fundamental with wavelength $4R$ and its harmonics.

For the time being, assume for instance that the field under consideration is a vector field which is not yet quantized. In the cavity and in its fundamental mode which amplitude takes the well known spherical standing wave form $A \frac{\sin(\pi r/R)}{r}$, we know that this amplitude is rather constant for $r < R_0$ if $R_0 \ll R$, which justifies our treatment of the previous section where we assumed a constant ρ inside the sphere of radius R_0 .

The total integrated energy of our standing wave bounded by R is simply some-

thing like

$$m_0 = A^2 \int_0^R \frac{\sin^2(\pi r/R)}{r^2} 4\pi r^2 dr \quad (25)$$

This is the bubble rest energy or in other words, its mass m_0 . But while the wave is an extended object much beyond R_0 and hence so is its energy everywhere inside the bubble of radius R , we located and computed the gravific effect of m_0 only in a very smaller sphere of radius R_0 at the center of the bubble, almost point like. So inside the volume delimited by the discontinuity our standing wave is now submitted to its own static gravitational field with gravific point mass m_0 . Assuming in a first order approximation that the wave function is not too much deformed by this potential, it is easy to compute the total self gravitational potential energy E_p of this spherical standing wave in the bubble. Delimited by the discontinuous potential barrier in its own central static gravitational field of its gravific point-mass m_0 , it's E_p is given by:

$$E_p = Gm_0 A^2 \int_0^R \frac{1}{r} \frac{\sin^2(\pi r/R)}{r^2} 4\pi r^2 dr \quad (26)$$

We can of course multiply and divide the latter expression by m_0 and get

$$E_p = m_0 \frac{\int_0^R \frac{-Gm_0}{r} \frac{\sin^2(\pi r/R)}{r^2} 4\pi r^2 dr}{\int_0^R \frac{\sin^2(\pi r/R)}{r^2} 4\pi r^2 dr} = -\frac{2.43Gm_0^2}{R} \quad (27)$$

With the exact expression for the gravitational potential $\frac{c^2}{4}(e^{-4\frac{Gm_0}{rc^2}} - 1)$ which in the weak field approximation gives back the Newtonian potential $-Gm_0/r$ we have just used to compute E_p , E_p gets modified

$$E_p = m_0 \frac{\int_0^R \frac{c^2}{4}(e^{-4\frac{R_0}{r}} - 1) \frac{\sin^2(\pi r/R)}{r^2} 4\pi r^2 dr}{\int_0^R \frac{\sin^2(\pi r/R)}{r^2} 4\pi r^2 dr} \quad (28)$$

We numerically compute

$$E_p / \left(\frac{-Gm_0^2}{R} \right) = \frac{R}{R_0} \frac{\int_0^R \frac{1}{4}(1 - e^{-4\frac{R_0}{r}}) \sin^2(\pi r/R) dr}{\int_0^R \sin^2(\pi r/R) dr} \quad (29)$$

where we have introduced $R_0 = \frac{Gm_0}{c^2}$, and find that this expression fastly converges to 2.43765 when R/R_0 tends to infinity. For R smaller than R_0 the expression would instead decrease as R .

As a consequence, the self gravitationnal potential energy of our spherical wave in the bubble is a quantum inversely proportionnal to the bubble radius R : $E = h_G c / 4R$ provided we define $h_G = 4(2.43) \frac{Gm_0^2}{c}$: the gravitational Planck constant. Thanks to $c/4R = c/\lambda \approx \nu$, the frequency of our standing wave, we indeed obtain $E = h_G \nu$, the fundamental relation of Quantum Mechanics and by the way establish a direct link between the gravitational Planck constant and m_0 .

Of course m_0 is a fundamental mass of nature which is the same whatever the bubble radius R : in other words the bubble can expand or collapse at constant

m_0 . Only the gravitationnal self energy varies and vanishes for infinite d in which case for the corresponding center of isotropy the spin 1 field is spread all over the universe but is only observable as a gravific mass m_0 at $r=0$. This is actually just the point mass we postulated in the previous section where we did not contemplate self gravitating energies.

Notice that $c/d \approx \nu$ is needed but $c/d = \nu$ is not possible otherwise the wave could not be trapped by the spherical discontinuity. Hence the spin 1 field we considered inside our cavity could not be an exactly massless field however its mass term is allowed to be very small if the discontinuity is a huge one.

Also notice that we take m_0 to be the gravific mass in order to compute the self gravitating energy E_p but of course from the outside world point of view the self gravitational potential energy does contribute to the gravific mass which is now $m_0 + E_p$ and reduces to m_0 only in case the discontinuity is at infinity.

Now for any propagating wave packet of any kind of field (massive or not, charged or not, arbitrary spin, composite or not, for instance even the wave function of an as big molecule as a C60 fullerene for which the Quantum Mechanics predictions have been recently tested with flying colors) we can always consider its decomposition into spherical waves components. Each of these spherical waves with frequency ν can be created or annihilated by the bubble state we have just described provided the bubble center is located at the center of isotropy of the spherical wave and of course provided other quantum numbers are conserved in the overall process. The important result is that a quantum of such isotropic centripetal wave with frequency ν can be absorbed (annihilated from the point of view of the outside world) when a bubble which discontinuity was initially at infinity, instantaneously collapses and stabilizes at distance d allowing a standing wave frequency which is also ν inside the bubble. In this way a quantum of energy $E = h_G \nu$ from the initial wave is absorbed. If another identical quantum is absorbed the bubble will become twice smaller and so on.

For each quantum absorption or release, the increase or decrease of the self gravitational potential energy of the bubble and its size follow accordingly and change the bubble gravific mass from m_0 to $m_0 + E_p$ and back to m_0 . As we anticipated in a previous section, such mechanism allows gravific mass to be defined only near the considered center of isotropy as long as the wave is trapped. In between the network points where various fields are propagating, gravific mass is not defined. This is our understanding of the link between $T^{\mu\nu}$ and $S^{\mu\nu}$ which also provides an interesting picture to understand the origin of annihilation and creation operators of Quantum Field Theory.

Notice however our new prediction: the Planck constant should no more be a constant but should start to decrease about the high energy threshold corresponding to a wavelength about r_0 .

23. Quantum gravity in the solar system

Angular momentum must be quantized in the solar system according :

$$mvr = n\hbar_G$$

Considering that the protoplanetary disk had constant surface density the mass found between radius r and $r+dr$ is $m(r) \propto r$ while the fundamental relation of dynamics yields

$$v(r) \propto r^{-1/2}$$

if the gravitationnal field of our central protostar was dominant. Thus the disk must have fragmented and concentrated at quantified radii r_n :

$$r_n^{3/2} \propto n\hbar$$

The rotation period T is a quantity proportionnal to $r^{3/2}$ and must therefore be quantized. It was indeed shown by JM Souriau that solar system periods are not only approximate multiples of 30 days, but also occupy a Fibonacci suite of frequencies, in such a way that neighbour planets are minimally resonant. After fragmentation of the primordial disk, protoplanetary masses grew in such a way that $m(r) \propto r$ was lost while radii and periods remained in the same proportions. If so $\hbar_G$ can be estimated. Given that the protoplanetary disk radius was approximately the present solar system radius r_{Oort} and its mass 10^{-3} solar Mass, its initial surface density was

$$\sigma = \frac{M_\odot}{1000\pi (r_{Oort})^2}$$

and

$$\sigma 2\pi r_n (r_{n+1} - r_n) \frac{2\pi}{T_n} (r_n)^2 = n\hbar_G$$

If 30 days is the fundamental period then Venus and Earth are at $n=11$ and $n=12$ respectively but may be the smaller self-rotation period in the solar system, 10 hours for Jupiter, is a more likely fundamental one in which case $n_{venus} \approx 990$ and $n_{earth} \approx 1080$ (L. Nottale formula has a 3 days fundamental period).

$$\hbar_G \approx \frac{4\pi}{1000} \frac{M_\odot}{1000T_{venus}} \frac{(r_{earth} - r_{venus})(r_{venus})^3}{(r_{Oort})^2} = 6.10^{29} Js$$

to be compared with the Planck constant of electromagnetism

$$\hbar_q = 10^{-34} Js$$

It follows that

$$m_{vacuum} = \sqrt{\frac{\hbar_G c}{G}} \sim 10^{24} kg$$

of the order of one fifth of the earth mass.

$$d_{vacuum} \approx \frac{Gm_{vacuum}}{c^2} 1.7 \sim 1mm$$

Can vacuum effects related to this network of masses be tested? All laboratory experiments involve masses moving at about 300 km/sec relative to our vacuum masses provided these are at rest with respect to the CMB frame. Being alternatively attracted and repelled a free mass test should vibrate or be submitted to deformations with typical frequencies of the order of 10 MHz in vacuum. Anyway it is probably difficult to extract such signal from the noise since it affects in almost the same way the experimental setup. No doubt that the optimal conditions are those of free motion with highly reduce noise i.e. free fall in space. Gravity Probe B is a free falling apparatus having an extremely good control of deformation and motion of its gyroscopes (the most spherical ever man-made objects), rotating at 0.03 mm from their stator, and a read-out system highly torque sensitive which should render it optimal for the detection of our vacuum effects. It appears that indeed the experiment has discovered unexpected new phenomena among which resonance peaks in the drift rate of the gyroscopes axis.

If we were to adopt a more conservative point of view , we would have a single fundamental Planck constant for both gravity and electromagnetism and $m_{vacuum} \sim 3.10^{-8}kg$ and $d_{vacuum} \sim 3.10^{-35}m$.

24. Quantizing electromagnetism

The same method used to quantize gravity allows us to quantize electromagnetism with another Planck constant h_Q .

$$\frac{h_Q c}{d_e} = \frac{q^2}{4\pi\epsilon_0 d_e} \Rightarrow h_Q = \frac{q^2}{4\pi\epsilon_0 c} = \frac{e^2}{c} \quad (30)$$

assuming that in the same bubble we have in addition a genuine positronium electromagnetically bound state. But now the vanishing energy condition should apply by compensating the electromagnetic potential energy of the pair by its spin-spin magnetic energy.

$$\left(\frac{q\hbar_Q}{2m}\right)^2 \frac{1}{c^2 4\pi\epsilon_0 d_e^3} = \frac{q^2}{4\pi\epsilon_0 d_e} \quad (31)$$

Yielding the bubble radius $d_e = \frac{\hbar_Q}{2mc} = \frac{137}{2}r_e \approx 2.10^{-13}m$ where r_e is the classical radius of the electron ($2.8 \cdot 10^{-15}m$).

Motivated by the approximate $z_{CMB} = 1000 \approx 2\pi.137$ in

$$h_Q = z \cdot \frac{e^2}{c} \approx 2\pi.137 \cdot \frac{e^2}{c}$$

we suspect cosmological expansion to be responsible for a coevolution of $\alpha = \frac{e^2}{\hbar c}$ and the masses scale.

25. Gravity, Quantum mechanics and Spirituality

For several decades and in spite of theoreticians sustained efforts, quantizing gravity has raised the major issue, not solved to date, of the compatibility between

the conceptual foundations of General Relativity and Quantum Mechanics, the two pillars of contemporary physics. Indeed, these appear radically antinomic, the main obstacle at the origin of this incompatibility between MQ and GR certainly being the inexistence of any privileged coordinate system and in particular the impossibility to define an absolute time in GR. It is already encouraging to realize that such kind of obstacle immediately disappears within the framework of a theory as DG which is built starting from an absolute and non dynamical flat space-time, a familiar framework for quantization. But well beyond, it is not only the simple perspective of unification between two ways of thinking nor merely a quantization program that opens up with DG. Indeed the theory, appears to generate quantization from its own principles and does much more than throwing some new light on the well known interpretation issues of QM: it solves them for the most part.

In DG we find two cohabiting modes of gravity: a continuous source one having propagating wave solutions and a discrete source one, a network of point masses structuring the vacuum, each point being able to communicate via instantaneous gravity with all the others and having in its neighborhood and being the center of isotropy of a stationary wave system oscillating inside a finite volume delimited by a gravity discontinuity. As we have already shown, each such system can emit a spherical centrifugal wave or absorb a spherical centripetal wave thereby a packet of energy (quantum) proportional to the wave frequency. The fundamental relation of quantization linking energy and frequency of the absorbed/emitted wave, $E = h\nu$, is therefore a consequence of the theory. On the other hand, the network points, while absorbing and emitting in a non local and concerted way a new system of spherical waves, is perfectly able to trigger the collapse of any QM wave packet.

We can now reconsider the most important interpretational issues of QM and explain which kind of solution DG offers in each case. We advise the reader to first read the first ten pages of "The transactional interpretation of Quantum Mechanics" by JG Cramer where the seven issues (Identity, Complexity, Collapse, Non locality, Completeness, Predictivity and the uncertainty Principle) are introduced and discussed within the Copenhagen Interpretation.

- Identity

What is the state vector (or wave function) of QM? The wave packet collapse is a so enigmatic and unacceptable process for most physicists that a positivist interpretation which does not take serious the physical reality of the wave function eventually stood out, interpretation according to which it is no more than a tool for efficiently computing relations between observables. At the contrary we believe that QM waves are as real and on the same footing as classical electrodynamics wave solutions of Maxwell equations. The state vector thus describes a purely wave phenomenon propagating in the continuous space-time of DG. Only when detection occurs (interaction or measurement) and the wave packet collapses, a very different physical process than the propagating one, does the more localised particle aspect

manifests itself. This way of thinking is not new actually: it is the a priori most obvious way first considered and studied by de Broglie and Heisenberg then criticized and unfortunately abandoned due to the non locality issues this approach raises.

- Non Locality

The main reason why the collapse of the wave packet is so disturbing is that it is essentially non local. This is not only a prediction of the QM formalism but now an experimental fact after many historical experimental results (in particular the A Aspect one) have firmly established the existence of QM non local correlations in entangled systems. Thus one must recognize the strong physical reality of this process. DG allows to go one step beyond in the acceptation and visualisation of the process. The discrete mode of vacuum, the points network, by annihilating or creating a system of spherical waves, can trigger the collapse of any wave packet. This collapse is allowed to be non local since all points can communicate via DG instantaneous gravity.

- The Wave Packet Collapse (Why?, How?)

The collapse has to be concerted ("decision taken in common by all involved network points") in order to respect the Born probability law: the energy of the vibration at each space-time point determines the probability for the wave packet to collapse there. But there is no need for the transactional interpretation of J.G Cramer to justify this point. It simply results from the fact that for instance light intensity at each point is according classical electrodynamics given by the signal energy there, i.e the mean of the squared signal, i.e. the squared modulus of the complex amplitudes sum that enter in the composition of this signal. For what concerns a light beam which photons are emitted one after the other, in between the emitter points and receiver points we have nothing else but a wave packet $s(x,y,z,t)$, with energy at x,y,z always given by the temporal mean of $s^2(x,y,z,t)$. But absorption or energy emission (involved in detecting a photon) can only occur in quantum packets ($E = h\nu$) by the network points. Naturally we then expect that an energy absorption will be more probable when the available energy at a given point that the wave brings there is more important on the mean (at the particular time "chosen by the point networks" for the collapse during one period, if the instantaneous amplitude is not enough to provide the minimum energy at a given point the quantum will not be absorbed there). Therefore it is not surprising that the energy determines at least the mean probability that the wave packet collapse takes place at that point. ACTually the collapse is not only possible but mandatory because of the discrete way network points can absorb or emit energy. The particle aspect is only manifested in the collapse: there is no more necessity for the obscure if not paradoxical wave-particle duality of the Copenhagen interpretation in our framework, since the wave and particle aspects are not really dual

aspects of the same reality: they now just stand for the influence of the two independent DG modes of vacuum! The wave packet propagating and spreading in the continuous mode of vacuum space is just from time to time transformed into a new more localized wave-packet by the non local and concerted action of the discrete mode of vacuum.

- Complexity

Advanced waves of the transactional interpretation (positive energies going backward in time or negative energies going forward in time) are not available having been rejected from the formalism of modern Quantum Field Theory since at a second quantization level these are completely understood in terms of annihilation operators. The Born probability law thus can not be interpreted as a transaction between a retarded field and its complex conjugate advanced field. By the way, let us recall that antiparticles are not advanced waves since when they go backward in time, following the Feynmann point of view, they are negative energy objects (see www.darksideofgravity.com/antimatiere.htm).

- Predictivity

Maybe could we hope to be able to compute more than just a probability if we had access to more than a temporal mean of $s(t)$, its instantaneous value or mean on the reduced time interval where the collapse decision is taken by the point networks. It would allow hopefully to eliminate much of the indeterminism. If there is a hidden determinism that makes appeal to blind physical processes (a non spiritualist understanding), it is totally unknown and remains to be explored.

- The Uncertainty Principle

We know from Fourier analysis that the better the space-time localisation of a signal the poorer its localisation in the space of frequencies. The time-frequency principle of uncertainty is therefore purely classical and not a mystery. Only when energy is substituted to frequency thanks to the quantization relation in the uncertainty principle do interpretational issues arise. In our framework, the time frequency uncertainty principle comes with the wave physics that takes place in the continuous mode of vacuum. The discrete mode of vacuum on the other hand establishes the link between energy and frequency, so that we can derive immediately the energy frequency uncertainty principle that only deals with the detected quantum and the recreation of a new more localized wave packet. The same for all other uncertainty relations. Interpretational issues most often related to the obscure concept of duality are avoided in this way (see previous paragraphs)

Should the QM formalism evolve, leading to new possible testable effects? Certainly if more determinism is hidden. Even at the new level of understanding implied by DG, for the collapse to be possible by the discrete vacuum mode probably the usual formalism must already be modified since the spherical waves base restricted

to waves having as isotropic centers the network points is complete and considerably reduced compared to the more usual one which isotropy centers scanned the whole continuum.

There is still an important issue : what triggers and when the wave paquet collapse by the network ? We should not neglect the firmly spiritualist way of understanding motivated by a deep analogy that we find between the vacuum network and a more familiar one: the neural network of our branes. In the same way as a correspondance exists between the activity states of billion neurons in a brain and mental or consciousness states, in the same way the states (vibration modes) of all points in the vacuum cosmic network could represent the physical manifestation of a spirit or consciousness of the universe, the living mode of vacuum, the one which eventually triggers the collapse of all wave paquets and by the way the periodic reactualisation of the universe. The inter-subjective if not objective character of reality for all individual minds would be insured in this way. The individual minds could be those of all living beings in the universe, may be as many components of the larger and encompassing cosmic one, the neural network being in interaction, in a way that remains to be studied, with the global vacuum network in the volume occupied by our brane.

Cerebral neural networks play the role of a fundamental interface if their activation is an essential step in the process that leads to the collapse of wave paquets for instance if they provide the global network with the information (our brains would be the senses of the universe) necessary for it to decide the way the wave paquets should collapse may be by introducing in the process a certain level of determinism hence ordering and favouring in a discreet but efficient way some states among those that the probability amplitude gives equiprobables. Metaphysical outlooks are fascinating, particularly the idea that our of states of consciousness are shared by the global spirit which in turn could enlarge our faculties (if we are willing to morally improve ourselves) and give our intuition an access to an infinite bank of knowledge through modified consciousness states. Mind survival to the brain death and integration into the universal mind is one of the most fascinating possibility in this perspective.

26. Conclusion

We could settle down here the foundations of the new theory of gravitation on flat spacetime which necessarily imposes itself as soon as we give up the GR geometrical point of view. Eventually, we find that this allows to solve many long lasting theoretical issues such as negative masses and stability, QFT vacuum divergences and the cosmological constant, negative energy and tachyonic representations of the Lorentz Group but also leads to very remarkable predictions: Locally, the disagreement with GR only arises at the PPN level in the preferred frame of our isometries, black holes disappear and gravitomagnetism can arise in an unusual way. Globally, a softly accelerating necessarily spatially flat universe in good agreement with the

present data is a natural outcome of the model. The growing of primordial fluctuations works well in this dark gravity theory without any need for dark energy nor dark matter components and the context is very promising to help solving the galaxy or cluster of galaxies missing mass issues. At last, we derived a gravitational wave solution leading to the observed decay of the binary pulsar orbital period. More speculative but very promising developments have started to be explored in Ref 3). Interesting ideas related at some level to the ones explored here can be found in Refs. 69 68 70 71 72 75 (see also many references of interest therein).

References

1. JP. Petit, *Science. Astr. and Sp. Sc.* **226**, 273-307 (1995).
2. <http://www.jp-petit.com/science/f300/a301.htm>
3. <http://www.darksideofgravity.com/DG.htm>
4. T. Damour and I I. Kogan, hep-th/0206042.
5. S. L. Robertson, gr-qc/0502088.
6. F. Henry-Couannier, gr-qc/0404110.
7. F. Henry-Couannier, gr-qc/0410055.
8. S. Chen, hep-th/0203230.
9. S. Weinberg, *Gravitation and Cosmology* (John Wiley and sons, New York, 1972).
10. S. Weinberg, *Quantum Field Theory, Vol 1*
11. A. D. Linde, *Rept. Prog. Phys.* **47**, 925 (1984).
12. R. R. Caldwell, *Phys. Rev. Lett.* **91**, 071301 (2003).
13. P. H. Frampton, *Mod. Phys. Lett.* **A19**, 801 (2004).
14. S. M. Carrol, M. Hoffman and M. Trodden *Phys. Rev.* **D68**, 023509 (2003).
15. A. D. Sakharov, *JETP Lett.* **5**, 24 (1967).
16. A. D. Linde, *Phys. Lett.* **B200**, 272 (1988).
17. J. D. Anderson, *Phys. Rev.* **D65**, 082004 (2002).
18. M. Tegmark, gr-qc/9702052
19. G. Feinberg, *Phys. Rev.* **D17**, No 6.
20. E. Recami, arXiv:physics/0101108.
21. C. M. Will, Theory and experiment in gravitational physics. CUP
22. Piazza and Marinoni, *Phys.Rev.Lett.* 91 141301, astro-ph/0304228.
23. D.N. Spergel et al. (WMAP Collaboration)
24. W.T. Hu, astro-ph/9508126
25. A.C. Fabian et Al, astro-ph/0306036
26. N. Straumann, astro-ph/0006423
27. S. Deser, and J. Franklin gr-qc/0408067
28. J. Ehlers, et Al gr-qc/0510041
29. S. M. Kopeikin gr-qc/0105060
30. N. Rosen, A bi-metric theory of gravitation *J. Gen. Rel. and Grav.* 4,435-47
gr-qc/0101025
31. Y. Aharonov, and D. Bohm, Significance of Electromagnetic Potentials in the Quantum Theory *Phys. Rev.* 115, 3 (1959).
32. hep-th/9510023
33. www.cosmosafr.iap.fr/MIT-RG1F.htm
34. C. M. Will gr-qc/0510072
35. M. Milgrom *Astrophys. J.* 270, 365; 270,371; 270,384 (1983).
36. J. D. Bekenstein *Phys. Rev.* D70 083509 (2004)
37. R. Foot, H. Lew and R.R. Volkas *Phys. Lett.* B272, 67 (1991)
38. T. D. Lee and C.N. Yang *Phys.Rev.* 104, 256 (1956)
39. J. Brian Pitts and W. C. Schieve Slightly Bimetric Gravitation gr-qc/0101058
40. J. Brian Pitts and W. C. Schieve Null Cones in Lorentz-Covariant General Relativity
gr-qc/0111004
41. The Supernova Legacy Survey: Measurement of $\omega(m)$, $\omega(\lambda)$ and w from the first year data set. astro-ph/0510447
42. M. Carrera and D. Giulini On the influence of the global cosmological expansion on the local dynamics in the solar system gr-qc/0602098
43. <http://chandra.harvard.edu/photo/2006/1e0657/>
44. S. Hossenfelder Cosmological consequences of Anti-gravitation gr-qc/0605083

45. K. Trencovski E G. Celakoska On the Pioneer anomaly gr-qc/0604068
46. K. Tangen Could the Pioneer anomaly have a gravitational origin gr-qc/0602089
47. R. Schodel, T. Ott, R. Genzel Stellar dynamics in the central arcseconf of our galaxy astro-ph/0306214
48. <http://www.jp-petit.com/science/f700/f701.htm>
49. <http://www.darksideofgravity.com/quantumgravity.pdf>
50. P. Moyassari, S. Jalalzadeh IJMPA, Vol. 20, No. 11 (2005) 2515-2521
51. JP. Luminet, A. Riazuelo, R. Lehoucq, J. Weeks astro-ph/0312312 (2006)
52. W. de Boer, C. Sander, V. Zhukov, A.V. Gladyshev, D.I. Kazakhov astro-ph/0602325
53. A. Blanchard, M. Douspis, M. Rowan-Robinson, S. Sarkar astro-ph/0512085
54. D. Kirkman et al, Astrophys. J. Supp. **149**, 1 (2003)
55. B.D Fields and S. Sarkar astro-ph/0601514
56. <http://www.darksideofgravity.com/masses.htm>
57. <http://www.god-does-not-play-dice.net/download.html>
58. <http://www.god-does-not-play-dice.net/index1.html>
59. R. D. Sadykhov physics/0606240
60. M. Sethi, A. Batra, D. Lohiya astro-ph/9903084
61. M. Kaplinghat, astro-ph/9911066
62. F. Henry-Couannier et al., gr-qc/0507065
63. T. Schucker, I. ZouZou astro-ph/0511521
64. T. Schucker, A. Tilquin astro-ph/0506457
65. The Nearby SN Factory astro-ph/0401513
66. C.Rovelli hep-th/0310077
67. S. Gehlaut, P. Kumar, G. and D. Lohiya astro-ph/0306448
68. G. 't Hooft S. Nobbenhuis gr-qc/0602076
69. J.W. Moffat hep-th/0610162
70. C.C. Barros Jr. physics/0509011
71. G. Chardin Antimatter in general relativity? DAPNIA-04-57, Mar 2004.
72. L. Iorio gr-qc/0606092
73. L. Iorio gr-qc/0712-1273
74. L. Bergstrom, J. Edsjo, M. Gustafsson astro-ph/0602632
75. I. Bogdanoff G. Bogdanoff Topological field theory of the initial singularity of space-time. Class.Quant.Grav.18:4341-4372,2001.
76. Jin He astro-ph/0512614
77. Public communications with Glen Deen and Polestar101 in forum <http://www.physicsforums.com/showthread.php?t=104694>
78. A. Hajian astro-ph/0702723
79. F Henry-Couannier <http://www.darksideofgravity.com/armagfr.htm>
80. T Buchert J Larena J.M Alimi gr-qc/0606020
81. J.D Bekenstein astro-ph/0701848
82. Private communication with JM Virey, to be published.
83. M-T. Jaekel, S.Reynaud gr-qc/0502007
84. G. Esposito-Farese gr-qc/0705.4043
85. G.Lochak, L. Urutskoev, Low energy nuclear reactions and the leptonic monopole
86. E.H. Lewis, Microscopic Ball Lightning, Sciencejunk.org
87. K.Shoulders and S.Shoulders, Charge Clusters in Action
88. N. Bevis, M. Hindmarsh, M. Kunz, J. Urrestilla astro-ph/9911066
89. R. Aurich, H. S. Janzer, S. Lustig, F. Steiner astro-ph/0708.1420v2
90. D.J. Eisenstein et all astro-ph/0501171v1
91. A. Riazuelo astro-ph/0804.2437v1

92. L. Rudnick, S. Brown, L.R. Williams astro-ph/0704.0908
93. A. Kashlinsky, F. Atrio-Barandela, D. Kocevski, H. Ebeling astro-ph/0809.3734
94. Hao. Liu, Ti-Pei Li, astro-ph.CO/0907.2731
95. Hao. Liu, Ti-Pei Li, astro-ph.CO/1003.1073
96. C.L. Bennett et All, astro-ph.CO/1001.4758
97. I. Quiros, gr-qc/0411064
98. I.T. Drummond, astro-ph/0008234
99. E.Kh. Akhmedov, hep-th/0204048
100. N. Bevis, M. Hindmarsh, M. Kunz, J. Urrestilla astro-ph/0702223
101. U.Sawangwit et al , astro-ph.CO/0911.1352
102. <http://www.ifa.hawaii.edu/cosmowave/supervoids/>
103. B.Y. Welsh, R. Lallement, J.L Vergely, S.Raimond New 3D gas density maps of NaI and CaII interstellar absorption within 300pc, in AandA, Vol 510 02/2010



